

Fam. CETONIIDÆ.Subfam. *ISCHNOSTOMINÆ*.**ISCHNOSCELIS.**

Ischnoscelis, Burmeister, Handb. der Ent. iii. p. 179 (1842); Lacordaire, Gen. Col. iii. p. 489.

This genus, interesting as displaying an affinity with the Goliathinæ group of Africa and Asia, so far at least as the cornuted head of the males is concerned, seems peculiar to Mexico. Two species have been described, widely dissimilar in size, colour, length, and clothing of the male fore legs, and to a lesser extent in the armature of the head, but closely agreeing in all essential points of structure.

1. *Ischnoscelis hopfneri*.

Goliathus hoepfneri, Gory & Percheron, Monogr. Cétoin. p. 154, t. 26. fig. 2¹.

Ischnoscelis höpfneri, Burm. Handb. der Ent. iii. p. 182²; Westw. Linnæa Entom. x. p. 326, t. 2.

Hab. MEXICO^{1 2} (*Sturm, in coll. Sallé*).

There are two examples only, a male and a female, of this rare species in the Sallé collection.

2. *Ischnoscelis dohrni*.

Ischnoscelis dohrnii, Westw. Linnæa Ent. x. p. 326, t. 2¹.

Hab. MEXICO¹, Sonora (*in coll. Bates*).

Subfam. *GYMNETINÆ*.**ARGYRIPA.**

Argyripa, Thomson, Typi Cetonid. p. 11 (1878).

Allorhina, pars, Sallé, Bates (olim), Westwood.

Three species of this beautiful genus have been described, remarkable for the curious armature of the head and thorax in the males. The group appears to be restricted to Central America and the neighbouring North-western parts of South America.

1. *Argyripa lansbergei*.

Allorhina lansbergei, Sallé, Ann. Soc. Ent. Fr. 1857, p. 617, t. 13. no. iii. figg. 1, 1 a-c.¹

Allorhina hypoglauca, Westw. Trans. Ent. Soc. Lond. 1874, p. 475, t. 7. fig. 6².

Argyripa lansbergei, Thomson, Typi Cetonid. p. 11.

Hab. NICARAGUA².—COLOMBIA¹.

This species is not contained in our collections and does not appear to have been