

9. **Gymnetis chontalensis.** (Tab. XXIII. fig. 11.)

Gymnetis chontalensis, O. Janson, Cist. Entom. i. p. 374, t. 9. fig. 4¹.

Hab. NICARAGUA, Chontales (*Janson*¹, *Belt*).

In this species the sternal process is of remarkable vertical width, and rather short and abruptly bent downwards.

10. **Gymnetis coturnix.** (Tab. XXIII. fig. 12, var.)

Gymnetis coturnix, Burm. Handb. der Ent. iii. p. 289¹.

Hab. PANAMA, Chiriqui (*Ribbe*, coll. *Bates*).—SOUTH AMERICA, Colombia¹.

Var. *G. nigrina*. Griseo-nigra, confluerter nigro-maculata, thorace strigis nonnullis guttisque elytrorum marginalibus versus apicem carneis.

Hab. PANAMA, Chiriqui (*Trötsch*), Volcan de Chiriqui (*Champion*).

An example obtained from the late Edwin Brown's collection, and ticketed "Brazil," agrees with Burmeister's description of this species, and I suspect the locality attached is erroneous, the real habitat being most likely Colombia. A Chiriqui specimen from Ribbe agrees with it, except that the black spots of the elytra unite in larger patches. The ground-colour of this (the typical form) is above dark greyish or tawny-brown, with two or three distinct orange-coloured spots on the margin of the elytra. The species is extremely close to *G. chontalensis*. Besides the very different colour and the fewer and larger black spots of the elytra, I find no differences except that in *G. coturnix* the thorax has more acute and produced hind angles, and the black tubercle of the epimera is surrounded by a pink ring. The sternal process is a little less abruptly bent downwards, but the degree of this may be a variable character; it has the same great vertical width as in *G. chontalensis*. The male is of a uniform dark grey colour beneath as in *G. chontalensis*.

11. **Gymnetis balzarica.**

Gymnetis balzarica, O. Janson, Cist. Entom. ii. p. 580¹.

Hab. COSTA RICA (*Van Patten*).—ECUADOR, Balzar Mts.¹

One example of this well-marked species was obtained from Van Patten's collection. *G. balzarica* was previously known only from the Balzar Mountains on the Pacific side of the main Andes in Ecuador.

12. **Gymnetis chevrolati.** (Tab. XXI. fig. 16, var. *ramulosa*.)

Gymnetis Chevrolat, Gory & Perch. Monogr. Cétoin. p. 346, t. 69. fig. 3.

Gymnetis ramulosa, Bates, Trans. Ent. Soc. Lond. 1869, p. 389¹.

Hab. NICARAGUA, Chontales (*Belt*¹); PANAMA, Chiriqui (*Trötsch*), Volcan de Chiriqui 2000 to 3000 feet (*Champion*).