

IV.

15. **Euphoria chontalensis.** (Tab. XXIII. fig. 10.)

E. morosæ (Jans.) proxime affinis, sed differt inter alia thorace scutelloque politis nigro-æneis. Oblongo-quadrata, nigro-ænea, elytris solum opacis maculis plurimis transversis undulatis fulvis; supra capite densius, elytris sparsius, fulvo-setosa; subtus hirsutie sat rigida fulva; capite densissime punctato, clypeo prope apicem angustato apiceque alte reflexo nec sinuato; thorace disco sparsius, lateribus densius arcuato-punctatis, basi medio lævi, lobo basali sat fortiter emarginato; elytris utrinque costis duabus validis impunctatis, interstitiis distanter lineatim punctatis, apice suturali haud producto; pygidio sat dense setoso, crebre striguloso, nigro-æneo, maculis utrinque fulvis (interdum deficientibus). Processus sternalis brevis et latus, antice rotundatus. Tibiæ anticæ ♂ acute tridentatæ, duobus apicalibus approximatis et exstantibus, 3° valde remoto. Antennæ ♂ clava elongata, cæteris articulis conjunctis vix brevior.

Long. 17 millim.

Hab. NICARAGUA, Chontales (*Belt*).

Five examples, apparently all males.

16. **Euphoria precaria.**

Euphoria precaria, O. Janson, Cist. Entom. ii. p. 583¹.

Hab. PANAMA, Chiriqui (*Trötsch*), Bugaba, Tolé (*Champion*).—SOUTH AMERICA, Colombia¹.

Closely allied to *E. iridescens*, Schaum. One of our examples agrees with Janson's description with regard to the elytra being of a dark olive-green ground-colour (like the rest of the body), with some small obscure yellowish spots on the elytra; but in others there is as much dark yellow spotty-colouring on the elytra as dark green, and the sides of the elytra especially are yellow with a series of large, quadrate, but irregular, dark green spots. The thorax is dark green, with a fine dorsal line and a small basal spot on each side yellow; but in one example the thorax has four dark-green vittæ as in *E. iridescens*.

17. **Euphoria iridescens.** (Tab. XXII. fig. 1.)

Euphoria iridescens, Schaum, Analecta Entomologica, p. 45¹; Burm. Handb. der Ent. iii. p. 388.

Hab. MEXICO, Puebla, Izucar (*Sallé*), Guerrero (*Baron*); GUATEMALA¹ (*Sallé*), Mirandilla 1700 feet, Aceytuno (*Champion*).

V.

18. **Euphoria leprosa.**

Euphoria leprosa, Burm. Handb. der Ent. iii. p. 379¹.

Hab. MEXICO¹, San Andres Tuxtla (*Sallé*), Atoyac in Vera Cruz (*H. H. Smith*).

The form of the clypeus is intermediate between that of Section IV. and that of Section VII. There are two varieties of this species in our small series:—

1. Smaller and narrower, dull black, with three narrow chalky-white vittæ on the