

24. *Euphoria nigriventris*.

E. vestitæ affinis et similariter colorata, sed differt statura minore, magis quadrata, subdepressa, thorace medio basi vix emarginata, ♂ antennarum clava mediocri, sed articulis 1°-7^m conspicue longioribus tibiisque anticis sat latis tridentatis. Subtus nigro-nitida pedibus concoloribus; clypeo antice minime vel haud dilatato, fere quadrato, margine apicali medio paullo sinuato; thorace obtuse trapezoideo, medio paullo rotundatim dilatato, deinde usque ad basin lateribus paullulum sinuatis, opaco, dense piloso, fulvo, vitta lata utrinque discoidali (intus recta extus dilacerata) nigro-viridi; scutello opaco, impunctato; elytris obtuse costatis, subtilissime punctulato-striatis, fulvis, nigro-viridi (plerumque transversim) maculatis; pygidio dense confluentur striguloso.

Long. 11-12 millim. ♂.

Hab. MEXICO, Peras, Etla (*Sallé*), Jalapa (*Höge*).

The more typical examples are from Jalapa. Two from Etla and Peras are distinctly more elongate and are darker, the broad dull greenish-black thoracic vittæ invading nearly the whole surface. Eight examples, males.

25. *Euphoria quadricollis*.

E. nigriventri maxime affinis; differt magis elongata, thorace ante medium angulatim dilatato deinde ad basin lateribus parallelis, medio basi sicut in *E. nigriventri* minime emarginato; quoad colores vix differt, ventre rufo-nitido excepto. ♂. Antennarum clava articulis 1°-7^m paullo longior; tibiæ anticæ angustæ, sicut in *E. geminata*, dentibus duobus parvis prope apicem, 3° superiore obsoleto.

Long. 12 millim. ♂.

Hab. MEXICO, Durango city (*Höge*).

Two male examples, perfectly similar.

26. *Euphoria atra*.

Quoad formam *E. nigriventri* simillima, sed toto atra, supra opaca, subtus nitida, dense griseo-pubescent; thorace medio basi distincte sed minus quam in *E. vestita* sinuato; elytris fortiter costatis. ♂. Antennarum clava articulis 1°-7^m paullo longior; tibiæ anticæ tridentatæ, dente 3° minore acuto, remoto.

Long. 12 millim. ♂.

Hab. MEXICO, Peras (*Sallé*), Oaxaca (*Höge*).

Three examples, all males.

27. *Euphoria vestita*.

Cetonia vestita, Gory & Perch. Monogr. Cétoin. p. 271, t. 52. fig. 6¹.

Cetonia irregularis, Gory & Perch. loc. cit. p. 272, t. 53. fig. 1² (nec Olivier).

Erirhipis vestita, Burm. Handb. der Ent. iii. p. 387³.

Hab. MEXICO^{1 2 3}, Oaxaca, Capulalpam, Peras (*Sallé*), Jalapa (*Höge*), Cuernavaca (*H. H. Smith*); COSTA RICA, Cache (*Rogers*).

Burmeister describes this species as having, in the male, an excessively large antennal club and nearly simple anterior tibiæ. A few male examples answering to this description, with females corresponding in size and colour, have been received from the above-mentioned localities. The figures cited from Gory and Percheron represent