

NELEUS (p. 14).

Neleus tlascala (p. 14).

To the localities given, add:—HONDURAS, Ruatan Island (*Gaumer*).

Neleus interstitialis (p. 15).

To the localities given, add:—MEXICO, Teapa in Tabasco, Atoyac in Vera Cruz (*H. H. Smith*), Temax in North Yucatan (*Gaumer*), Tabi in Yucatan (*F. D. G.*).

RHODOCANTHOPUS (p. 15).

Rhodocanthopus maillei (p. 15).

To the localities given, add:—MEXICO, Omilteme and Amula in Guerrero, Teapa in Tabasco (*H. H. Smith*).

PTICHOPUS (p. 17).

Ptichopus angulatus (p. 17).

To the localities given, add:—MEXICO, Dos Arroyos in Guerrero (*H. H. Smith*), Temax in North Yucatan (*Gaumer*).

UNDULIFER (p. 18).

Undulifer incisus (p. 18).

To the Mexican localities given, add:—Omilteme in Guerrero (*H. H. Smith*).

SORANUS (p. 19).

9. Soranus depressifrons.

S. recticorni similis, sed elytris relative longioribus. Sat anguste elongatus, politus; labrum antice paullo angustatum, sinuato-truncatum, grosse punctatum; clypeus latus, lævis, margine subrecto; frons toto depressa, medio transversim concava, piloso-punctata, cornu frontali nullo carinisque brevissimis; thorax relative parvus, antice distincte angustatus, subtiliter punctulatus, juxta marginem anticum fere sicut in gen. *Veturio* paullo prominens et convexus, sulco marginali antice lato et profunde fere usque lineam dorsalem extenso, lateribus impunctatis foveaque transverse lævi; elytra relative elongata, dorso subdepressa, punctato-sulcata; prothorax subtus tibiæque dense fulvo-hirsuta.

Long. 25 millim.

Hab. MEXICO, Huitzilac in Morelos (*Höge*). Two examples.

The leaflets of the antennal club are short (not more than twice as long as the joints), but the species does not well fit into any of the genera of the restricted group 'Passalini' which have this character. The prominence of the anterior margin of the thorax in the middle, limited on each side by the end of the deep and broad marginal sulcus, is a character of the genus *Veturius*; it is, however, only slight in the present case, and