

oblique spot extending from the suture to the margin. The prosternal chin-piece is gently and widely emarginate. The prosternum is closely and finely rugose, slightly pubescent in the middle; the intercoxal process is very slightly convex, truncate at the apex. The metasternum is closely punctured, with an elongate central puncture behind. The abdomen is rather finely punctured at the apex, rugose at the base; all the middle part of the basal segment clothed with yellowish velvety pile. The sides of the second segment are clothed with yellow pile, as are also the posterior angle of the metasternum and the whole of the posterior coxæ. The lateral carina of the basal segment of the abdomen is deeply angularly excised about the middle, with a yellow spot above.

5. *Paradomorphus emarginatus*.

Elongatus, angustus, nitidus; capite cyaneo-viride, longitudinaliter sulcato; thorace cyaneo, obsolete transversim striolato, tenue haud crebre punctato; elytris purpureis, crebre subasperato-punctatis, postice attenuatis, apice profunde emarginato, vitta suturali maculaque ante medium pallide flavis.

Long. 3 lin.

Hab. MEXICO, Acapulco in Guerrero (*Höge*).

The head is moderately closely and finely punctured; the longitudinal channel is rather deep at the top of the face, shallow in front; the face is rather flat, with a very slight ovate swelling on each side of the median channel. The thorax is somewhat narrowed at the base; shining, the striation not very distinct; transversely impressed at the base, with a shallow round fovea in the middle of the posterior margin; the posterior ridge is distinct, curved outwards and downwards to meet the margin beyond the middle. The elytra have the subcosta very distinct, gradually approaching the suture posteriorly; each elytron has a small pale yellow spot at about one third from the base, and a narrow sutural stripe extending from rather behind the middle to the apex; the apex itself is very deeply cut out, leaving the sutural angle acutely denticiform. The prosternal chin-piece is arcuate, nearly straight in the middle. The prosternum is strongly punctured, sparsely pubescent; the intercoxal process is subparallel, with the lateral margins thickened, raised, and smooth, the apex truncate, with the angles slightly prominent. The metasternum has a small elongate central puncture, with an elongate white spot at some distance from the side. The abdomen is finely and not very closely punctured; the basal segment with a white stripe at some distance from the side; the second and third segments each with a small white spot at the side.

6. *Paradomorphus elegantulus*.

Cupreo-æneus, parum nitidus, capitis vertice thoraceque purpureis, elytris nigro-æneis flavo-maculatis; capite antice crebre rugoso, vertice strigoso; thorace transversim strioloso, postice transversim leviter impresso, ad angulos anticos macula alba notato; elytris creberrime subasperato-punctatis, aureo-maculatis, apice emarginato, denticulato.

Long. $2\frac{3}{4}$ lin.