

steel-blue; the lateral carina of the basal segment has a very slight sinuosity in the middle.

10. *Paradomorphus corrugatus*.

Æneus, elytris obscure rufo-cupreis; capite crebre rugoso, antice convexiusculo; thorace antice bene convexo, fortiter rugoso, punctato; elytris depressis, crebre asperatis, regione suturali obsolete parce flavo-griseo-pubescente; corpore subtus parce brevissime flavo-griseo-pubescente.

Long. 3-3½ lin.

Hab. MEXICO, Juquila, Toxpam (*Sallé*).

The head in some examples is finely, in others coarsely, rugose; this is no doubt a sexual distinction. The thorax is very convex; the lateral impression is not deep, and does not extend to the anterior angles (it is scarcely visible from above), and in some lights there is an indication of a longitudinal median impression; the posterior ridge is very distinct, curved. The elytra have the subcosta only very slightly marked in the female, more distinct in the male; the apex is acuminate, finely serrate. The prosternal chin-piece is gently arcuate. The prosternum is closely and not very finely rugose; the intercoxal process is rather flat and pubescent, very slightly narrowed posteriorly, emarginate at the apex. The metasternum has a fine impressed median line; the episterna are finely punctured and pubescent. The abdomen is moderately closely punctured, the basal segment at the sides and at the intercoxal process very closely so; the latter is obtusely angular. The lateral carina of the basal segment is rather strongly and widely sinuate about the middle.

11. *Paradomorphus persimilis*.

Præcedenti similis et affinis, differt tamen elytris postice magis acuminatis; processu prosternali ad apicem emarginato, angulis paullo elevatis.

Long. 3-3½ lin.

Hab. MEXICO, Chilpancingo in Guerrero (*Höge*).

I separate this species with doubt from the preceding. It only differs in having a more decided longitudinal impression on the thorax. The elytra are rather more acuminate, and have the subcosta very distinct. The prosternum is more closely punctured, and the angles at the apex are slightly raised.

12. *Paradomorphus complexus*.

Subdepressus, æneo-cuprascens, nitidus, thorace postice elytrisque olivaceo-virescentibus; capite sat crebre punctato, medio canaliculato-impresso; thorace parum convexo, punctis transversis crebre aspersis, medio leviter bi-impresso; elytris creberrime asperatis, apice parum serrulato.

Long. 4½ lin.

Hab. NICARAGUA, Chontales (*Belt*).

The head has the punctures fine, but distinct, and slightly separated from each other; the median longitudinal impression is slightly narrowed at the middle of the