

post medium ampliatis, dein ante apicem angustatis, ad apicem paullo divaricatis, maculis plurimis irregularibus æneis confertim subtiliter rugulosis et griseo-pilosis impressis, interstitiis glabris, nitidis.

Long. 5 lin.

*Hab.* NICARAGUA, Chontales (*Belt*).

The head is strongly and closely punctured, except on the inner side of the round swelling on each side of the lower part of the face; there is a very slight prominence above each eye. The thorax has its anterior margin somewhat raised in the middle, and projecting very slightly over the head; the impressions are marked with numerous short, curved, or wavy deeply impressed lines; the sides are rugulose; on each side of the central impression there is a small smooth spot on the elevated part. The elytra are transversely impressed at the base (leaving the shoulders somewhat elevated), and gradually curved down to the apex; the costa on each is very obtuse, interrupted at intervals by the small impressions; there are strong punctures on and just below the shoulder, and at the apex; and at the side behind the middle there are numerous asperities. The prosternal process is very strongly and closely punctured. The middle of the short metasternum and the intercoxal process of the abdomen are sparsely and delicately punctured. The sides of the abdomen are clothed with fine pubescence of a slightly golden colour, which forms a large spot on the second segment and a small one on the first and third segments.

## 2. *Agrilus diversus*. (Tab. IV. figg. 11, 11 a.)

Nigro-purpureus; capite cupreo, crebre punctato, fronte supra incisa, binodosa; thorace crebre punctato et flexuoso-strigoso, disco sat profunde impresso, lateribus rugulosis, antice oblique impressis, ad angulos posticos carina brevi lævi; elytris confertim subtiliter rugulosis, dorsim planatis, postice inclinatis, pallide griseo-maculatis, apice denticulato, lateribus obscure æneo-viridibus; prosterno creberrime punctato.

Long. 5 lin.

*Hab.* MEXICO, Toxpam (*Sallé*); GUATEMALA, Chacoj in Vera Paz (*Champion*).

The ovate punctured swellings on the face of this species are on the upper part raised above the level of the eyes. The thorax has its discoidal impression not so deep as in the preceding species, narrow in the middle, widest behind; the raised parts on the disc are generally tinted with green. The pale grey pubescence which forms the markings on the elytra is extremely short. Each elytron has a small acute tooth at the apex near the suture. The punctuation on the prosternum is very dense, less so on the metasternum (though confluent transversely). The abdomen in the middle is less closely and much less strongly punctured; the apical segments sparsely clothed with pale pubescence. The middle of the metasternum and abdomen are æneous. One example has the sides of the metasternum slightly coppery.

## 3. *Agrilus oculatus*.

Fusco-cupreus, parum nitidus; capite fortiter excavato; thorace fortiter punctato et vermiculoso, inæquali; elytris rugosis (maculis irregularibus lævibus notatis), fusco et æneo variegatis, litteris guttisque griseo-pubescentibus ornatis.

Long. 3-3½ lin.