

9. *Agrilus distans*. (Tab. V. figg. 2, 2 a, b.)

Viridi-æneus, subtus brunneo-æneus, sat nitidus, creberrime asperato-punctatus; elytris maculis sex albo-pubescentibus.

Long. 4 lin.

Hab. MEXICO, Oaxaca (*Höge*).

The head has some closely placed converging curved punctured lines on the vertex; the face is rather strongly and closely punctured; there is a very shallow impression in the middle. The thorax has two very shallow inconspicuous impressions on the disc, one anterior, the other posterior; the punctures are moderately close together, transversely elongate, and often united to each other transversely by fine lines; on each side of the front of the disc the surface is rather more rugulose; the usual lateral impression is brownish-æneous, with some whitish pubescence at the anterior part; the posterior ridge is distinct, smooth—commencing from the angle and curving round, it nearly touches the side about the middle. The elytra are closely and somewhat strongly asperate-punctate; the subcosta is well marked and sculptured in the same way as the rest of the surface; the basal impression is well marked and slightly pubescent; just before the middle there is a spot close to the suture, and at some distance from the apex another more elongate and whiter spot. The prosternum is very closely and rather strongly punctured; the episterna are closely punctured and clothed with pale grey pubescence. The metathoracic episterna are rather closely and moderately finely asperate-punctate. The abdomen is distinctly and not very closely punctured. The lateral carina of the basal segment is distinctly angulated rather behind the middle, with a small spot above (fig. 2 b).

10. *Agrilus squaliformis*.

Agrilus squaliformis, Thomson, Typi Bupr. p. 85¹.

Hab. MEXICO, Cordova, Puebla (*Sallé*); GUATEMALA¹, Dueñas, San Gerónimo 3000 feet (*Champion*), Chimaltenango (*Conradt*).

This species has the lateral carina of the basal segment of the abdomen very strongly angulated behind the middle.

Three examples, two from Mexico and one from Guatemala, are of a beautiful steel-blue colour instead of the usual brassy-æneous.

11. *Agrilus divaricatus*. (Tab. IV. figg. 16, 16 a, b.)

Brunneo-æneus, subtus cuprescens, nitidus; capite antice longitudinaliter sulcato, macula flavo-pilosa ornato; thorace crebre subtiliter punctulato, transversim aciculato, disco medio sat profunde sulcato; elytris postice bene attenuatis, ad apicem parum divaricatis et cupreis, regione suturali impresso et creberrime punctato; abdomine medio quam elytra latiore, supra utrinque macula magna flavo-pilosa ornato; episternis metathoracis flavo-pilosis.

Long. 6 lin.

Hab. MEXICO, Cuernavaca in Morelos (*Sallé*, *Höge*), Almolonga, Iguala in Guerrero (*Höge*).