

24. *Agrilus excisus*. (Tab. V. figg. 6, 6 a.)

Obscure brunneo-æneus, parum nitidus, subtus cuprascens; capite antice rufo-cupreo, subopaco; thorace sat crebre punctato et transversim ruguloso, disco bi-impresso, angulis posticis linea nitida curvata instructis; elytris ruguloso-punctatis, maculis octo pallide griseis ornatis, ad apicem cupreis, paullo ampliatis, singulo exciso, denticulato.

Long. $3\frac{2}{3}$ lin.

Hab. PANAMA, Bugaba 800 to 1500 feet (*Champion*).

The head is longitudinally impressed on the forehead, which is bigibbose; the punctures on the front are moderately close together and not very distinct, those on the vertex being much stronger and longitudinally confluent. The thorax has two shallow impressions on the disc, one at the base, and one in front of the middle; the posterior ridge is curved and punctured, and consequently not very conspicuous. The elytra have the punctures rather strong and united transversely by double parallel lines; the pale whitish-grey spots are placed in the sutural impression—one in the basal impression, one before the middle, one behind the middle, and one smaller one near the apex; at the apex they are parallel; each is emarginate, denticulate at the outer angle and sutural angle, with a strong acute spine at the outer side of the emargination (fig. 6 a). The prosternum is very closely and moderately strongly punctured, slightly shining, sparsely pubescent. The metathoracic episterna are moderately closely and strongly punctured and sparsely pubescent. The abdomen is shining, very finely and moderately closely punctured; at the sides are two obscure sparsely pubescent pale greyish spots; the lateral carina of the basal segment alters its direction posteriorly (fig. 6).

25. *Agrilus cribricollis*. (Tab. V. figg. 8, 8 a-c.)

Omnino brunneo-æneus, parum nitidus; thoracis disco fortiter sat crebre punctato, transversim ruguloso, medio in longitudinem profunde impresso; elytris tenuiter flavo-griseo-pubescentibus, regione suturali creberrime subtiliter punctulato, maculis nonnullis fere calvis nitidis ornatis, lateribus confertim rugosis, apice ipso rufo-cupreo, denticulato.

Long. $4\frac{1}{2}$ lin.

Hab. MEXICO, Jalapa (*Höge*).

The head is coarsely punctured on the vertex, triangularly impressed in front, the face very finely punctured and pubescent, with two small smooth raised spots in the middle; the clypeus and a line running along the inner side of the eye shining coppery. The thorax is coarsely punctured on the disc, with the intervals confluent transversely in front; the deep longitudinal impression is indistinctly punctured, narrowed in front; the sides are finely rugose; the posterior ridge is well defined, shining, rather swollen and raised anteriorly (fig. 8 a). The elytra are sparingly clothed with short greyish-yellow pubescence, except at the apex (fig. 8 c). Each elytron has four bare, more shining spots—the first, near the base, does not touch the subcosta nor the suture; the second and third touch the subcosta but not the suture;