

the epipleuræ, yellowish-testaceous; the legs fusco-piceous, the basal two thirds of the femora reddish-testaceous.

Hab. GUATEMALA (*Sallé*). One female example.

Var. γ. Above and beneath testaceous, the head piceous-brown, the prothorax, legs, and antennæ (except towards the apex) reddish-testaceous.

Hab. GUATEMALA, Las Mercedes (*Champion*). One male example.

This species varies very much in size and colour. It is chiefly distinguished from its allies by its coarsely punctured head and thorax, long antennæ in the male, thickly and confusedly punctured elytral interstices, and hairy legs. *S. corrosa* in its elytral sculpture and hairy legs resembles *S. brevipilis*; but it differs from that species in its less elongate shape, more prominent eyes, more slender antennæ, shorter thorax, different colour, &c.

54. *Statira horrida*. (Tab. II. fig. 16, ♀.)

Elongate, rather convex, subopaque, blackish-violaceous, the prothorax reddish-testaceous, the body above and beneath thickly clothed with moderately long erect ashy hairs. Head coarsely and rugosely punctured, shallowly grooved on each side between the eyes anteriorly, the latter moderately large; antennæ rather stout, moderately long, piceous, the apical joint (♀) not so long as joints 8–10 united; prothorax convex, about as long as broad, the sides immarginate, rounded anteriorly, and strongly constricted behind, the hind angles moderately prominent, the basal margin grooved within, the surface coarsely and rather closely punctured; elytra with very shallow intrahumeral depression, rather long, subparallel to beyond the middle, very coarsely and deeply punctate-striate, the punctures very closely packed and strongly transverse to beyond the middle and thence to the apex crenate, the interstices quite narrow, moderately convex to about the middle, strongly so beyond, and finely and confusedly punctured, the apices obtuse; beneath piceous, shining, somewhat thickly punctured; legs rather long, piceous, thickly clothed with long erect ashy hairs.

Length $7\frac{1}{2}$ millim.; breadth 2 millim. (♀.)

Hab. GUATEMALA, Panzos in Vera Paz (*Champion*).

A single female example, captured by myself in the lower part of the Polochic valley. *S. horrida* resembles *S. corrosa*, the head and thorax being somewhat similarly shaped and punctured, but differs greatly in the sculpture of the elytra and also in the upper and under surfaces and legs being much more thickly clothed with long erect hairs. The rows of punctures on the basal half of the elytra are closely packed and very strongly transverse, and the interstices in consequence quite narrow, the latter being confusedly punctured.

55. *Statira punctatissima*.

Moderately elongate, rather dull, brownish-piceous, thickly clothed with short erect hairs, the prothorax and the front of the head testaceous, the elytra subopaque and with the shoulders and the suture (narrowly) testaceous. Head coarsely and closely punctured, the eyes moderately large; antennæ (♂) long, fusco-ferruginous, the basal joint testaceous, the apical joint nearly as long as joints 7–10 united; prothorax convex, about as long as broad, the sides immarginate, rounded anteriorly, and strongly constricted behind, the hind angles moderately prominent, the basal margin raised, the surface very coarsely, deeply,