

16. **Eugnampthus rufifrons**, sp. n. (Tab. II. fig. 7.)

Longulus, parce setosus, niger, elytris cyaneis, vertice rufo; elytris subtiliter seriatim punctatis.
Long. cum rostro $4\frac{1}{4}$ millim.

Hab. MEXICO, Totosinapam (*Sallé*).

Antennæ black, very slender, with the long club rather less slender, each joint of the club narrowed towards the base, the terminal joint much shorter than the tenth. Rostrum short, as long as the head, the eyes very prominent, the head almost impunctate. Thorax very slender, but slightly conical, sparingly subobsoletely punctate, shining. Elytra very elongate, with very regular series of fine punctures, the interstices broad, flat, without any distinct punctuation. Legs very slender, the middle tibiæ of the male strongly curved. In the female the eyes are rather smaller and more distant.

One of the most slenderly-formed members of the genus. It was labelled in Sallé's collection *E. rufifrons*, Chevr. Three examples.

17. **Eugnampthus niger**, sp. n.

Niger, nitidus, griseo-setosus, elytris nigro-subcyaneus; prothorace parce punctato; elytris seriatim crebre sat fortiter punctatis, interstitiis modice angustis, haud omnino planis.
Long. cum rostro $4\frac{1}{4}$ millim.

Hab. MEXICO, Ventanas and Refugio in Durango (*Höge*).

Antennæ moderately long, the club not linear, each joint being narrowed towards the base, the terminal joint quite acuminate, shorter than the tenth. Rostrum rather longer than the head; eyes large, approximate to the neck, the space between them finely punctate at the sides, smooth and shining on the middle. Thorax rather slender, moderately finely and not closely punctate. Elytra blue-black, shining, the punctures in the series rather closely placed. Legs very slender.

Allied to *E. rufifrons*, but independently of the colour of the head it differs in numerous details. We have received only two specimens, which I presume are males, the apex of the middle tibia being slightly produced internally and acuminate.

18. **Eugnampthus longulus**, sp. n.

Longulus, cyaneus, griseo-setosus, nitidus, antennis pedibusque nigris; prothorace gracili, parce, haud subtiliter sed levissime punctato; elytris elongatis, seriatim crebre sat fortiter punctatis, interstitiis modice angustis, haud omnino planis.
Long. cum rostro 5–6 millim.

Hab. MEXICO, Parada, Juquila (*Sallé*), Oaxaca (*Höge*).

Closely allied to *E. niger*, but more elongate in form, with the surface of a beautiful blue colour, and with the punctures of the thorax so slightly impressed that they are indefinite and the surface irregular. The four examples received include both sexes, the male having the middle tibia incurved and strongly acuminate at the inner angle.