

**19. *Eugnamptus semivittatus*, sp. n.**

Testaceus, setosus, elytris longitudinaliter nigro-signatis, antennis fuscis, apice flavo; elytris crebre fortiter profundeque seriatim punctatis.

Long. cum rostro 4 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

Rostrum about as long as the thorax. Club of the antennæ elongate and very slender, the terminal joint more than half as long as the tenth, the apical portion of the latter joint, as well as the following one, yellow. Head short, the eyes very large, the space between them almost impunctate. Thorax very slender, coarsely and rather closely punctate. Elytra yellow, with a black lateral stripe, and a second dark stripe near the suture, this inner stripe indefinitely prolonged beyond the middle and imperfectly connected with a black mark joining the outer stripe; the punctures of the series are large, very deep, and closely placed. Legs red, the hinder and middle tibiæ a little infusate at the apex externally. Two specimens.

**20. *Eugnamptus obscurus*, sp. n. (Tab. II. fig. 8.)**

Longulus, gracilis, niger, supra fuscus, opacus, elytris fusco-testaceis, antennis pedibusque fusco-rufis; capite prothoraceque omnino densissime punctatis, rugosis.

Long. cum rostro  $4\frac{1}{2}$  millim.

*Hab.* GUATEMALA, Panajachel 5000 feet (*Champion*).

Club of the antennæ slender, but with each of the joints narrowed towards the base, the joints preceding the club very long, the terminal joint only a little shorter than the tenth. Rostrum very short, shorter than the head, densely rugose. Eyes large, the space between them very densely rugose. Thorax slender, with excessively dense, coarse, rugose sculpture. Elytra with deep striation and rugose interstices, the intercalated series at the base of the first interstice unusually elongate. Legs long and slender.

Only one specimen has been obtained of this extremely distinct species; this was found upon the margin of the Lake of Atitlan.

**21. *Eugnamptus varius*, sp. n. (Tab. II. fig. 9.)**

Gracilis, subtus fusco-niger, nitidus, supra opacus, setosus, fuscescens; elytris irregulariter fusco testaceoque variegatis; antennis pedibusque testaceis, fusco-variegatis.

Long.  $3\frac{1}{4}$  millim.

*Hab.* GUATEMALA, Cerro Zunil (*Champion*).

Antennæ with moderately long club, each joint of which is narrowed to the base, the terminal two joints darker than the ninth, which is yellow. Rostrum minute, densely rugose. Eyes rather large, distant from the neck, the space between them excessively densely and coarsely punctured. Thorax slender, longer than broad, excessively densely and coarsely punctured. Elytra very deeply striate, the striæ densely and coarsely punctured. Legs yellow, vaguely marked with fuscous. The only speci-