

rather longer than the thorax in one sex, shorter in the other. Eyes not very large, rather widely separated. Head elongate. Thorax very narrow, subcylindric, closely, very coarsely punctate. Elytra slender, with very regular series of coarse deep punctures. Legs slender, clothed with erect hairs.

This insect has somewhat the appearance of the more slender of the species of *Deporaus*. It differs in many respects from *E. niger*, our only other black *Eugnamptus* without spots or marks. We have received only two mutilated examples.

25. ***Eugnamptus brevicollis***, sp. n. (Tab. II. fig. 11.)

Niger, griseo-setosus, elytris subpurpureo-tinctis, rostro, capite, prothorace femoribusque basi rufis; elytris densissime fortissimeque regulariter seriatim punctatis, interstitiis perangustis.

Long. cum rostro  $4\frac{1}{2}$  millim.

Var. rostro prothoracisque marginibus nigris.

Hab. PANAMA, Volcan de Chiriqui 3000 to 4000 feet (*Champion*).

Antennæ intensely black, pilose, the club very elongate, quite linear, the terminal joint a good deal shorter than the tenth. Rostrum very broad, quite short, the eyes moderately large; head sparingly punctate. Thorax short, about as broad as long, very coarsely, not densely punctate. Elytra with regular rows of very densely packed large and deep punctures. Legs black, the base of femora bright red, the tarsi obscure red.

We have received eight examples of this very distinct species.

26. ***Eugnamptus cornutus***, sp. n. (Tab. III. fig. 1.)

Elongatus, nigerrimus, prothorace læte rufo; setis erectis concoloribus parce vestitus; rostro (maris?) superne apicem versus incrassato, parte anteriore in cornu reclinato, apice furcato, producto.

Long. cum rostro 6 millim.

Hab. MEXICO, Chilpancingo in Guerrero 4600 feet (*H. H. Smith*).

Antennæ quite black, long, the club not very long. Head elongate, quite sparingly punctate. Thorax bright red, rounded at the sides, sparingly punctured. Elytra with quite regular series of coarse punctures, the interstices not punctate but setose, the first at the base with about seven intercalated punctures.

We have received only one example of this species; it is probably a male. The structure of the front of the rostrum is unique, but the curious horn is not conspicuous owing to its being depressed.

§ 2. *Pubescence of the body shorter, very fine, reclined.*

27. ***Eugnamptus salvini***, sp. n. (Tab. II. fig. 12.)

Brevior, testaceus, pubescens; capite prothoraceque densissime punctatis; elytris profunde lateque striatis, striis dense et fortiter punctatis.

Long. cum rostro  $3\frac{1}{2}$  millim.

Hab. GUATEMALA, Pantaleon, Mirandilla, Zapote, La Tinta (*Champion*).