

### 11. *Apion glyphicum*, sp. n.

Robustum, nigrum, supra parce subtiliusque albido-setosum, opacum, elytris subæneis; rostro robusto, capite inter oculos striato; prothorace fortiter punctato; elytris tenuiter minus profunde sulcatis, interstitiis planis.

Long.  $2\frac{1}{2}$  millim.

*Hab.* MEXICO, Jalapa (*Höge*).

There is but little difference between the sexes in this species; but the male has the middle and hind tibiæ mucronate at the apex, and the rostrum not quite so long as the head and thorax, while in the female it is about as long as the head and thorax. Rostrum stout, sulcate on each side; antennæ inserted near to the eyes, these latter convex, the interval separating them only moderately broad, and bearing three or four fine striæ. Thorax greatly narrowed from the base to the front, very coarsely punctate, with a very short channel in front of the scutellum. Elytra convex, much broader than the thorax; the grooves rather fine and not very deep, bearing distant, remote punctures, the intervals flat and rather broad. Legs stout. The pubescence of the under surface less scanty than that above, but still not close. Three specimens.

### 12. *Apion tabogense*, sp. n.

Breve, robustum, nigrum, subopacum, parcissime albido-setosum; rostro (maris) crassiusculo, rugoso; prothorace lato, conico, dense punctato; elytris striatis, interstitiis latis, planis.

Long. 2 millim.

*Hab.* PANAMA, Taboga Island (*Champion*).

We have received only two males of this species, but as it will be recognized without much difficulty—not being very near to any other—I have described it. The middle and hind tibiæ are mucronate at the apex, but the mucrones are slender spurs directed straight downwards, not inwards as in most of the other species. The rostrum is rather thick, curvate, densely sculptured except at the tip; the antennæ are inserted not very far from the eyes, being separated from them by not quite one fourth of the length of the rostrum; the eyes are rather large and prominent, placed close to the thorax, the interval between them moderately broad. The thorax is rather broad, much narrowed towards the front, scarcely at all impressed at the sides, moderately closely and coarsely but obsoletely punctate, with a small fovea in front of the scutellum. Elytra short and broad, coarsely striate, though the striæ are scarcely deep enough to be called sulci, the interstices rather broad, flat. The white setosity is not quite so scanty on the under surface as it is above, and under the orbits is a little more conspicuous.

### 13. *Apion championi*, sp. n. (Tab. III. fig. 4.)

Robustum, nigrum, sat nitidum, fere nudum; rostro medioeri, sculpturato, basi incrassato; prothorace conico, parce subtiliterque punctato; elytris convexis, striatis, interstitiis latis, planis.

Long.  $2\frac{3}{4}$  millim.

*Hab.* PANAMA, Volcan de Chiriqui, David, Caldera, Los Remedios (*Champion*).