

76. **Apion terminale**, sp. n.

Angustum, nigerrimum, nitidum, tenuiter albido-setosum, antennis pedibusque testaceis; rostro brevi, recto; elytris striatis, ante apicem impressis, apice subtumidulo.

Long. 2 millim.

Hab. MEXICO, Amula in Guerrero in August (*H. H. Smith*).

Closely allied to *A. longicolle*, but with the femora as well as the tibiæ and tarsi yellow; the thorax shorter, and the upper surface bearing a scanty scattered setosity, the unusual formation of the apices of the elytra more pronounced, and the sides of the breast quite covered with white setosity.

One example.

77. **Apion inflatipenne**, sp. n. (Tab. III. fig. 19.)

Breve, nigrum, nudum; elytris viridi-cyaneis, latissimis, ad basin quasi deplanatis, subtiliter sulcatis; pedibus nigris, femoribus in medio rufis.

Long. $2\frac{3}{4}$ – $3\frac{1}{2}$ millim.

Hab. GUATEMALA, El Reposo (*Champion*).

Rostrum rather stout, cylindrical, curved, longer than the head and thorax, punctate, shining; eyes somewhat convex; antennæ inserted a little behind the middle of the rostrum. Thorax rather short, greatly narrowed in front, somewhat finely but not densely punctate, with a short, deep, foveiform channel in front of the base in the middle. Elytra remarkably broad, appearing almost as if inflated, but somewhat deplanate in front, with rather fine, not deep sulci and very broad, flat interstices, somewhat dull. Under surface without any setosity. Middle coxæ well separated; hind legs black, with the larger part of the femora red.

Two specimens.

78. **Apion latipenne**, sp. n.

Breve, nigrum, nitidum; elytris viridi-cyaneis, latis, convexis, profunde minus late sulcatis; prothorace conico, parce punctato; femoribus late rufis.

Long. $2\frac{1}{2}$ millim.

Hab. PANAMA, Peña Blanca (*Champion*).

Rostrum rather short, cylindrical, shining, sparingly punctate, sulcate above on each side; antennæ inserted about one-third of the length from the base; eyes but little prominent. Thorax somewhat small, greatly narrowed in front, rather finely and sparingly punctured. Elytra very broad and short, very convex, much narrowed in front, with moderately deep and broad grooves. Legs not very stout; middle coxæ but little separated; femora red, except at the tips. One specimen.

Though similar in appearance to *A. inflatipenne*, this is, no doubt, distinct from that species, as it differs in numerous slight particulars, among others in having a very fine, minute, and scanty setosity on the thorax and elytra.