

- b*⁴. Elytral setæ shorter and decumbent *striatus*, Boh.
*b*³. Elytra with the alternate interstices interruptedly costate.
*c*⁴. Third interstice with two very large oblong tubercles; outer
 striæ coalescent; humeri less prominent *quadrigibbus*, sp. n.
*d*⁴. Third interstice with a single moderately large tubercle below
 the base; outer striæ free; humeri very prominent *ruptus*, sp. n.
*d*². Rostrum shorter, subquadrate; humeri prominent.
*c*³. Elytra with the alternate interstices more or less raised; outer
 striæ coalescent *grypsatus*, Boh.
*d*³. Elytra with the interstices feebly uniformly convex; outer striæ
 free *verapacis*, sp. n.
b. Wings rudimentary; scutellum smaller; anterior femora unarmed;
 humeri not prominent; rostrum subquadrate.
*c*¹. Elytra with the alternate interstices more or less interruptedly
 costate *chiriquensis*, sp. n.
*d*¹. Elytra with the alternate interstices more or less raised.
*e*². Elytral setæ broad, squamiform, raised. *subapterus*, sp. n.
*f*². Elytral setæ decumbent, narrow, hair-like *cinericius*, sp. n.

1. ***Eustylus sexguttatus***, sp. n. (Tab. XIII. fig. 20, ♂.)

Elongate, narrow (♂), broader (♀), piceous; densely clothed above with brown and whitish scales, the latter condensed into two vittæ on the disc of the prothorax and a stripe down each of the alternate elytral interstices, the elytra also each with three subquadrate blackish spots (two on the third, and one on the seventh interstice), and sometimes another small spot towards the apex; the scales on the under surface and along the flanks of the prothorax and elytra almost wholly whitish; the surface also set with minute scattered decumbent hairs. Rostrum longer than broad, widened anteriorly, the short deep scrobes converging posteriorly; eyes very large, separated by about their own width as seen from above; antennæ fully one-half the length of the body in ♂, the scape stout and reaching considerably beyond the front margin of the prothorax, joint 2 of the funiculus longer than 1. Prothorax slightly broader than long, narrowed and feebly constricted anteriorly, flattened down the middle of the disc, the punctures covered by the vestiture. Scutellum large. Elytra elongate-subtriangular in ♂, broader and gradually widened to the middle in ♀, the humeri prominent; punctate-striate, the alternate dorsal interstices costate. Anterior femora clavate and sharply dentate. Anterior and intermediate tibiæ strongly unguiculate in both sexes.

Length 7-10, breadth $2\frac{1}{5}$ - $3\frac{1}{2}$ millim. (♂ ♀.)

Hab. PANAMA (*Salvin*), Volcan de Chiriqui, David (*Champion*), Tabernilla, Canal Zone (*Busck*, in *U.S. Nat. Mus.*), Taboga I. (*Champion*, *Busck*).

Numerous examples, varying in the intensity of the markings, according to the predominance of the whitish or of the brown scales, the darker individuals having the white line on the fifth elytral interstice obsolete and the blackish spots very conspicuous, and the paler specimens the spots almost obsolete. *E. sexguttatus* may be at once separated from the Mexican *E. striatus* by the sharply toothed anterior femora, apart from its greater size, larger eyes, longer antennæ, shorter prothorax, and different elytral markings.