

2. *Eustylus setipennis*, sp. n. (Tab. XIII. figg. 21, 21 a, ♂.)

Elongate, narrow, rather convex, piceous; above densely clothed with brown scales, with cinereous or greenish-white scales intermixed, the lighter ones sometimes condensed into two faint vittæ on the disc of the prothorax and a more or less distinct short oblique streak on the outer part of the elytra at about the middle, the scales on the flanks in part, and those on the under surface, usually cinereous or greenish-white; the elytra also set with numerous long, stiff, suberect, blunt setæ down each of the dorsal interstices, the rest of the surface with short curled decumbent scattered setæ, the tibiæ pilose within. Head and rostrum canaliculate, the rostrum longer than broad, the scrobes subparallel as seen from above, the head flattened between the eyes and abruptly convex behind them, the eyes rounded and very prominent; antennal scape very stout, reaching beyond the front margin of the prothorax, joint 2 of the funiculus longer than 1. Prothorax much longer than broad, subcylindrical, slightly rounded at the sides, sparsely punctate. Scutellum small. Elytra moderately elongate, punctate-striate, the alternate dorsal interstices sometimes becoming more or less costate posteriorly, the humeri not very prominent. Legs stout, anterior femora unarmed.

Length $5-6\frac{1}{2}$, breadth $1\frac{1}{2}-2$ millim. (♂.)

Hab. GUATEMALA, Senahu and San Juan in Alta Vera Paz (*Champion*), Cacao near Trece Aguas (*Schwarz and Barber, in U.S. Nat. Mus.*).

Five examples, apparently all males, varying in the colour of the scales, the Senahu specimen having the flanks of the prothorax and elytra in great part greenish-white. Near *E. striatus*, Boh., differing from it in the subparallel scrobes, the longer antennal scape, and the strongly setose elytra.

3. *Eustylus striatus*. (Tab. XIII. fig. 22, ♂.)

Eustylus striatus, Boh. in Schönh. Gen. Curc. vii. 1, p. 42¹.

Length $5\frac{1}{2}-7$, breadth $1\frac{3}{4}-2\frac{1}{3}$ millim. (♂ ♀.)

Hab. MEXICO (*Hegewisch, in coll. Sommer*¹).

An elongate, narrow form, variegate or lineate above with a dense clothing of light and dark brown scales (the dark scales clustered into three narrow vittæ on the prothorax and various interrupted lines or small confluent spots on the elytra), and also set with numerous, rather long, curved, adpressed, dark setæ, the scales on the under surface greyish or opalescent; the short deep scrobes converging posteriorly; the eyes large and prominent; the antennal scape very stout and reaching the front of the prothorax; the prothorax elongate; the elytra very gradually narrowing from the base in ♂, broader, subparallel in their basal half, and more abruptly acuminate in ♀, the alternate interstices costate; the anterior femora unarmed. This species is not represented in any of the extensive Mexican collections received by us, but there are six examples of it in the Sommer collection, now in the Oxford University Museum, the male figured having been kindly presented by Prof. Poulton.

4. *Eustylus quadrigibbus*, sp. n. (Tab. XIII. figg. 23, 23 a, ♂.)

Moderately elongate, narrow (♂), broader (♀), piceous or ferruginous; densely clothed with coppery-brown scales, the prothorax with an indication of two faint paler stripes on the disc and the elytra each with an irregular, oblique, dirty white fascia just beyond the middle; the surface also set with scattered, curled, decumbent setæ. Head and rostrum canaliculate, the rostrum longer than broad, widened