

Species moderately large or small, elongate-ovate, oblong-ovate, or subovate, squamose, with the rostrum curved, the eyes large, distant or well-separated, and partly exposed (covered in *C. consobrinus*), the antennal club ovate, the prothorax bisinuate at the base and with the subtruncate hind angles received into the excavate base of the elytra, the alternate elytral interstices usually more or less raised, the mesosternum arcuately emarginate (horseshoe-shaped in *C. quadriplagiatus*, *C. carinifer*, and *C. evanescens*), the metathoracic episterna becoming very narrow anteriorly, the ventral segment 2 as long as, or longer than, 3, the femora unidentate or unarmed, the anterior tarsi sometimes (*C. bifenestratus*, *C. tirunculus*, and *C. cancellatus*) with a few projecting hairs in the ♂. (Nos. 62-88.)

62. **Cryptorrhynchus quadriplagiatus**, sp. n. (Tab. XXXII. figg. 31, 31 a, ♀.)

Oblong-ovate, black, the antennæ and the tips of the tarsi ferruginous; thickly clothed with brown and blackish scales, the prothorax with a broad white cruciform vitta on the disc (extending from the base to the apex), and the elytra each with a very broad, oblique, white fascia extending from a little below the shoulder to the suture, as well as an oblong, common, white apical patch (which nearly joins the fasciæ in front and is sinuous on its outer edge), the blackish scales clustered along each side of the prothoracic vitta and along the edges of the elytral markings; the upper surface also set with scattered short setæ, the legs annulate. Head densely, rugulosely punctate, the eyes somewhat distant; rostrum moderately stout, curved, about as long as the head and prothorax, rugulosely punctate and finely carinate at the base, for the rest minutely punctate, the antennæ inserted far behind the middle, joint 2 of the funiculus elongate, as long as 1, the club ovate. Prothorax broader than long, rounded at the sides, narrowed and constricted in front, deeply bisinuate at the base; densely, finely punctate, the disc apparently depressed down the middle and with an abbreviated median carina. Scutellum rounded. Elytra a little wider than the prothorax, oval, widening for a short distance below the base and narrowed thence to the apex; seriate-punctate, the interstices rugulose and sparsely granulate, 3, 5, 7, and 9 more or less raised, 3 becoming more prominent towards the base, the suture swollen posteriorly. Mesosternum horseshoe-shaped. Beneath finely punctate. Legs rather slender; femora sharply unidentate.

Length  $7\frac{1}{10}$ , breadth 3 millim. (♀.)

*Hab.* GUATEMALA, Cerro Zunil 4000 feet (*Champion*).

One specimen. Smaller and less elongate than *Diaporesis distincta*, the prothorax shorter and with a broad cruciform white patch on the disc, the three white patches on the elytra so enlarged as to become almost coalescent, the alternate dorsal interstices costate.

63. **Cryptorrhynchus carinifer**, sp. n. (Tab. XXXII. fig. 32.)

Oblong-ovate, black, the antennæ ferruginous; thickly squamose, the prothorax black, with a spot in front of the scutellum and the flanks anteriorly ochreous, the scutellum ochreous; the elytra with the base, a spot on the disc below this, and a large triangular patch at the sides towards the apex, black, the rest of their surface ochreous, except laterally and near the suture, where they are fulvous; the vestiture of the under surface and legs in great part ochreous, the femora and tibiæ fusco-annulate; the prothorax and elytra also set with short, scattered, rather coarse, semierect setæ. Head rugosely punctate, carinate down the middle, the eyes rather prominent and not very widely separated; rostrum strongly curved, moderately long, stout, rugosely punctate and sharply carinate at the base, and for the rest sparsely, very finely punctate, the antennæ inserted towards the base, joints 1 and 2 of the funiculus subequal in length, the club ovate. Prothorax broader than long, rounded at the sides anteriorly, narrowed and feebly constricted in front; coarsely, closely, shallowly punctate, and obsoletely carinate. Scutellum rounded.