

of the elytra. A single specimen from Guatemala apparently belongs to it, and also one from Mexico in the British Museum, the latter showing a pallid transverse space on the elytra beyond the dark basal patch, and therefore more like Herbst's figure of *C. obtentus*. The scattered, semierect, setiform scales on the elytra are coarse.

**66. *Cryptorrhynchus collinus*, sp. n.** (Tab. XXXIII. fig. 2, ♂.)

Elongate-ovate, shining, black, the antennæ and the tips of the tarsi obscure ferruginous; thickly clothed with blackish or brown scales, the prothorax with a narrow cruciform patch on the disc (sometimes reduced to a short streak at the base), the scutellum, and one or two small spots on the third elytral interstice at about the middle, cinereous or ochreous, the elytra also with scattered, irregularly distributed, short black streaks, the legs mottled or annulate with cinereous; the upper surface sparsely set with short, fine, suberect setæ; the vestiture of the under surface cinereous. Head rugulosely punctate, the eyes distant; rostrum stout, curved, not longer than the prothorax, rugosely punctate (except along the centre towards the apex), finely carinate at the base, much smoother in the ♀, the antennæ inserted at (♂) or behind (♀) the middle, joints 1 and 2 of the funiculus subequal in length, the club ovate, acuminate. Prothorax broader than long, rounded at the sides anteriorly, narrowed and strongly constricted in front; densely, somewhat coarsely punctate, the anterior lobe smoother, the posterior lobe sometimes with a short median carina. Scutellum rounded. Elytra moderately long, a little wider than the prothorax; seriate-punctate, the interstices punctulate and faintly granulate, 3 and 5 raised, the others moderately convex. Beneath closely punctate; first ventral segment depressed down the middle in the ♂. Mesosternum broad, prominent, feebly emarginate. Femora unidentate. Tibiæ almost straight, angulate at the base externally.

Length  $5\frac{1}{5}$ – $5\frac{3}{4}$ , breadth  $2\frac{1}{10}$ – $2\frac{1}{2}$  millim. (♂ ♀.)

*Hab.* GUATEMALA, El Tumbador, Pacific slope (*Champion*: ♂); PANAMA, Volcan de Chiriqui (*Champion*: ♀).

Three specimens, the male in very fresh condition. Near *C. obtentus*, but with the elytral vestiture much darker, and the black scales condensed into irregular, scattered, transverse or oblong streaks (instead of a large, common, subtriangular patch at the base), the setæ much finer.

**67. *Cryptorrhynchus divexus*, sp. n.** (Tab. XXXIII. fig. 3, ♀.)

Elongate-ovate, rather narrow, shining, black, the antennæ obscure ferruginous; thickly clothed with blackish scales, the elytra with a very large, common, transverse, indeterminate, cinereous patch a little beyond the middle, the legs broadly annulate with the same colour; the upper surface also set with short, scattered, setiform scales; the vestiture of the under surface cinereous. Head rugulosely punctate, the eyes distant; rostrum curved, of the same length as the prothorax, sparsely punctate, rugose and feebly carinate at the base, the antennæ inserted behind the middle, joints 1 and 2 of the funiculus equal in length, the club ovate, acuminate. Prothorax broader than long, rounded at the sides anteriorly, narrowed and strongly constricted in front; densely, somewhat coarsely punctate. Scutellum oval. Elytra a little wider than the prothorax, convex, elongate, narrowing from about the middle, the humeri obtuse; seriate-punctate, the interstices faintly granulate, 3 and 5 slightly raised, the others feebly convex. Beneath closely punctate. Mesosternum broad, prominent, feebly emarginate. Femora unarmed.

Length  $6\frac{1}{4}$ , breadth  $2\frac{1}{2}$  millim. (♀.)

*Hab.* GUATEMALA, Cerro Zunil (*Champion*).

One specimen. This species approaches *C. obtentus* and *C. collinus*, but it has more