

Three specimens. Less robust than *C. bioculatus*, the setæ much shorter and inconspicuous, the prothorax narrowed behind and without lighter markings, the whitish patch on the disc of each elytron transverse and smaller, the femora more feebly dentate. The following is yet another nearly allied form.

**85. *Cryptorrhynchus diophthalmus*, sp. n.** (Tab. XXXIII. fig. 20, ♂.)

More elongate than *C. distigma*, the scutellum and a transverse patch at about the middle of each elytron white; the upper surface somewhat thickly set with short, rather coarse, semierect setæ; the prothorax less constricted in front and more coarsely punctate; the elytra more coarsely seriate-punctate; the antennæ inserted towards the base in both sexes; the femora unarmed; the fifth ventral segment broadly truncate at the apex in the ♂.

Length 4, breadth  $1\frac{2}{3}$  millim. (♂ ♀.)

*Hab.* MEXICO, Jalapa (*Höge*), Toxpam (*Sallé*).

Two specimens, both from the Atlantic slope. The posterior femora are slightly angulate beneath, but they cannot be described as toothed. In one example there is a small white spot on the prothorax in front of the scutellum.

**86. *Cryptorrhynchus leucostigma*, sp. n.**

Oblong-ovate, nigro-piceous; clothed with rather coarse brown scales, the elytra variegated with black and also with a faint, interrupted, transverse white fascia before the middle; the upper surface somewhat thickly set with erect, pallid and dark setæ, the legs setose. Head densely punctate, the eyes distant; rostrum short, stout, feebly arcuate, finely punctate, rugulose at the base, the antennæ inserted behind the middle, joints 2-7 of the funiculus slender, 2 shorter than 1, the club ovate. Prothorax transverse, rather convex, rounded at the sides, narrowed and slightly constricted in front, feebly sinuate at the base; dull, densely, finely punctate. Scutellum small. Elytra very little wider than the prothorax, rapidly narrowing from about the middle; shining, rather coarsely seriate-punctate (the punctures not hidden by the vestiture), the interstices convex, flat along the suture. Beneath densely punctate. Mesosternum broad, prominent, arcuate-emarginate. Legs short; femora unidentate.

Length 3, breadth  $1\frac{1}{2}$  millim. (♀ ?)

*Hab.* GUATEMALA, San Juan in Vera Paz (*Champion*).

One specimen. In this obscure little species the elytra are very little wider than the prothorax, faintly fasciate with white, and rather coarsely seriate-punctate, the entire upper surface setose, the setæ nowhere clustered into fascicles. It approaches *C. degressus*.

**87. *Cryptorrhynchus leucaspis*, sp. n.**

Ovate, narrow, piceous, the antennæ ferruginous, densely clothed with intermixed pale brown and fuscous scales, the prothorax with a narrow pallid median vitta, which is whitish in front of the scutellum, the elytra with a common, broad, whitish space on the disc, extending from a little below the base to near the apex and widened in front and behind; the upper surface also set with short, coarse, erect setæ, the legs setose. Head densely punctate, the eyes rather small, laterally placed; rostrum short, stout, feebly curved, rugosely punctate to the tip, the antennæ inserted at the middle, joints 3-7 of the funiculus transverse and closely articulated, 2 short, the club ovate. Prothorax a little wider than long, narrowed and feebly constricted in front; densely punctate. Elytra punctate-striate, the interstices 3, 5, and 7