

Species small, oblong-ovate, squamose and setose, with the rostrum short, stout, widened towards the base and apex, the antennal club ovate, the eyes large, the prothorax produced and strongly fasciculate at the apex, with the ocular lobes feebly developed and the base bisinuate, the elytra subtriangular, produced at the apex, and with oblong setigerous prominences on the disc, the mesosternum prominent and feebly emarginate, the metathoracic episterna narrow, the ventral segment 2 longer than 3, the legs rather slender, the femora moderately clavate and unidentate. (No. 93.)

**93. *Cryptorrhynchus subcaudatus*, sp. n.** (Tab. XXXIII. figg. 25, 25 a.)

Subelliptic, rather narrow, shining, piceous, the antennæ, rostrum, and apices of the tarsi rufescent; thickly clothed with small, intermixed pale brown, fulvous, and cinereous scales; the prothorax with two fascicles of long, erect, dark brown setæ at the apex and a few shorter setæ scattered across the middle; the elytra also with scattered, semierect, light and dark setæ, the latter becoming crowded on the dorsal elevations. Head rugulosely punctate, somewhat swollen above the eyes, the latter moderately large and distant; rostrum short and stout, a little widened towards the base and apex, feebly curved, rugulosely punctate at the base and thence to the apex sparsely, minutely punctate, the antennæ inserted far behind the middle, joints 1 and 2 of the funiculus equal in length, the others short and widening outwards, the club stout, ovate. Prothorax about as long as broad, constricted and much narrowed in front; densely punctate, obsolete carinate at the middle. Scutellum small. Elytra much wider than the prothorax, subtriangular, strongly produced at the apex, the humeri slightly swollen; seriate-punctate, the interstices punctulate, 1 with an oblong setigerous prominence before, and 2 with a similar prominence at the middle, 3, 5, and 7 also raised near the base, 5 becoming convex posteriorly. Beneath closely, rather finely punctate. Mesosternum very prominent, feebly emarginate. Legs rather long; femora moderately clavate, unidentate.

Length  $4\frac{1}{2}$ , breadth 2 millim. (♀?)

*Hab.* PANAMA, Volcan de Chiriqui (*Champion*).

One specimen. This insect is very like *Anomæmerus tristis*, Faust \*, from Venezuela; but it has the rostrum broader towards the tip, and the antennæ are inserted behind the middle; the anterior femora are not nearly so stout; the elytra are strongly produced at the apex, and have the setigerous elevations differently placed, &c.

Species very small, ovate, sparsely and finely squamose, with the rostrum curved, the eyes large and well-separated, the antennal club ovate, the prothorax feebly bisinuate at the base and bifasciculate at the apex, the scutellum small, the elytra subtriangular, with the interstices narrowly raised, the third becoming more swollen towards the middle, the mesosternum very prominent and feebly emarginate, the metathoracic episterna broad, the ventral segment 2 longer than 3, the legs rather slender, the femora unidentate. (No. 94.)

**94. *Cryptorrhynchus guttatus*, sp. n.**

Ovate, piceous, the apical half of the rostrum, the antennæ, and tarsi more or less ferruginous; sparsely clothed with minute fulvous scales, the head with a small spot in front, and the elytra with a short streak beneath the humeri, a small spot at the middle of the second interstice, and another at its base, white or whitish, the prothorax with two fascicles of short, erect, fine, blackish setæ at the apex. Head

\* The type of this species is a male, and it has the anterior tarsi clothed with laterally projecting hairs.