

98. *Cryptorrhynchus opaciventris*, sp. n.

Oblong-ovate, piceous, the antennæ and the tips of the tarsi ferruginous; somewhat thickly clothed with small pale brown scales, with cinereous scales intermixed, the elytra mottled with black; the upper surface also set with erect setæ, the legs setose; the vestiture of the under surface sparse. Head densely punctate, the eyes moderately large and well-separated; rostrum short, feebly curved, slightly widened at the base, rugosely punctate to the tip in the ♂, much smoother in the ♀, the antennæ inserted at or a little behind the middle, joint 2 of the funiculus shorter than 1, the club ovate. Prothorax transverse, somewhat rounded at the sides, narrowed and feebly constricted in front; densely punctate. Scutellum small. Elytra considerably wider than the prothorax; punctate-striate, the interstices feebly convex, 3 slightly swollen at the base. Beneath opaque, alutaceous, finely punctate. Mesosternum raised, prominent, feebly emarginate. Legs short; femora acutely unidentate, the tooth on the anterior pair narrow; tarsi slender.

Length $3\frac{1}{2}$, breadth $1\frac{1}{2}$ millim. (♂ ♀.)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One pair. Near *C. simplex*, but smaller, the eyes not so large and more distant, the under surface opaque and alutaceous, the third elytral interstice slightly swollen at the base. The elytral striæ are less coarsely punctate than in *C. echinatus*, and the setæ are shorter and finer than in that insect.

99. *Cryptorrhynchus disjunctus*, sp. n.

Oblong-ovate, shining, piceous, the antennæ ferruginous; clothed with small brown scales, the prothorax with two curved blackish marks on the disc behind and a minute whitish spot in front of the scutellum, the latter ochreous, the elytra with a black spot on the third interstice below the base and a whitish one on the second beyond the middle; the upper surface also set with short, erect, light and dark setæ, the legs setose; the vestiture of the under surface sparse and pale. Head densely punctate, the eyes rather large and somewhat distant; rostrum short, stout, feebly curved, sparsely, minutely punctate, rugulose at the base, the antennæ inserted at about the middle, joint 2 of the funiculus much shorter than 1, the club ovate. Prothorax transverse, constricted and much narrowed in front; densely punctate. Elytra considerably wider than the prothorax, rapidly narrowing from about the middle; punctate-striate, the interstices punctulate, 3 swollen just below the base. Beneath rather closely punctate. Mesosternum prominent, arcuate-emarginate. Legs short; femora unidentate.

Length $4\frac{1}{4}$, breadth 2 millim. (♀.)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One specimen. Near *C. quadrituberculatus*, but with an oblong black prominence at the base of the third elytral interstice only (the one on the second interstice wanting), the second with a sharply-defined whitish spot beyond the middle. *C. guttatus* is marked in the same way, but that insect wants the black patch on the third elytral interstice, and has the second interstice swollen to near the middle, the prothorax binodose at the apex, &c.

100. *Cryptorrhynchus quadrituberculatus*, sp. n. (Tab. XXXIII. fig. 29, ♀.)

Oblong-ovate, shining, piceous, the antennæ and the tips of the tarsi ferruginous; somewhat thickly clothed with small fulvous scales, variegated with blackish and white scales, the latter condensed into two irregular, curved series of spots on the disc of the elytra, the blackish scales mainly clustered into two spots at the base of the prothorax and an oblong spot on the swollen portion of the second and third