

elytral interstices; the upper surface also set with long, erect setæ, the legs setose. Head densely punctate, the eyes rather large and well-separated; rostrum short, stout, feebly curved, finely punctate, rugulose at the base, the antennæ inserted at a little behind the middle, joints 3-7 of the funiculus short, 2 nearly as long as 1, the club ovate. Prothorax transverse, constricted and narrowed in front; densely, finely punctate. Scutellum oval. Elytra considerably wider than the prothorax, narrowing from a little below the base, produced at the apex; punctate-striate, the interstices feebly convex, 3 swollen near the base and 2 at the middle. Beneath dull, closely punctate. Mesosternum prominent, broad, feebly arcuate-emarginate. Legs rather short; femora subclavate, sharply unidentate; tarsi slender.

Length $2\frac{2}{3}$ - $3\frac{1}{2}$, breadth $1\frac{1}{8}$ - $1\frac{3}{4}$ millim. (♂ ♀.)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Four specimens, varying a good deal in size, the white spots on the elytra conspicuous in one only of them. The oblong, blackish elytral prominences—one near the base of the third interstice and the other at the middle of the second—are characteristic. The femora are more sharply dentate than in *C. opaciventris*.

101. *Cryptorrhynchus senticosus*, sp. n.

Oblong-ovate, shining, piceous, the elytra more or less mottled with rufo-piceous, the antennæ and the tips of the tarsi ferruginous; thickly clothed with brown scales, sometimes with a few whitish scales intermixed, the prothorax usually with two small patches at the base and the elytra with a few scattered spots or streaks black or blackish; the upper surface also set with rather coarse, erect, scattered setæ, the legs setose; the vestiture of the under surface sparse. Head densely punctate, the eyes moderately large and well-separated; rostrum about as long as the prothorax, curved, stout, sparsely punctate, rugulose and slightly widened at the base, the apical half smoother in the ♀, the antennæ inserted at a little behind the middle, the club ovate. Prothorax transverse, narrowed and feebly constricted in front; densely punctate. Scutellum small. Elytra considerably wider than the prothorax, rapidly narrowing from about the middle; punctate-striate, the interstices punctulate and feebly convex. Beneath rather sparsely punctate. Mesosternum prominent, arcuate-emarginate. Legs short; femora feebly unidentate; posterior tibiæ similarly formed in the two sexes, not excavate at the apex in the ♂.

Length 3- $3\frac{1}{2}$, breadth $1\frac{1}{4}$ - $1\frac{1}{2}$ millim. (♂ ♀.)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Ten specimens, in rather dirty condition. An obscure species, resembling *C. uncipes* in the colour of the vestiture, but smaller and narrower, the scutellum not so large, the legs shorter, the posterior tibiæ simple in the two sexes. The allied *C. simplex* is also larger, and has larger eyes, longer legs, and the vestiture somewhat differently coloured. The larger and more approximate eyes, the longer setæ, the non-vittate prothorax, the more sparsely punctured ventral surface, and the obsoletely dentate femora distinguish *C. senticosus* from *C. tectus*.

102. *Cryptorrhynchus contaminatus*, sp. n. (Tab. XXXIII. fig. 30, ♀.)

Oblong-ovate, shining, nigro-piceous, the antennæ and the tips of the tarsi ferruginous; the prothorax sparsely, the rest of the upper surface thickly, clothed with fulvous scales, the elytra with an oblique whitish streak on the disc below the base, a broad, angulate, common, post-median, whitish fascia, and some scattered black markings, both along the suture (before and beyond the post-median fascia) and on the disc, the scutellum whitish; the upper surface also set with rather long pallid and black setæ; the legs setulose, the femora subannulate. Head densely, rugulosely punctate, the eyes rather small, distant; rostrum feebly curved, not longer than the prothorax, widened, rugosely punctate, and subcarinate at the base, and for the rest