

narrowing thence to the apex; with rows of coarse, oblong punctures placed in shallow striæ, the interstices feebly convex and sparsely punctulate. Beneath sparsely, finely punctate. Mesosternum prominent, arcuate-emarginate, the walls declivous from the base. Legs short; femora feebly unidentate.

Length 5, breadth $2\frac{1}{10}$ millim. (♀.)

Hab. GUATEMALA, Cerro Zunil, Pacific slope (*Champion*).

One specimen, apparently somewhat worn. The dense white transverse patch on the head, the shining, sparsely squamose, and coarsely setose upper surface and legs, the rather long elytra, the sloping mesosternum, &c., sufficiently distinguish *C. phaleratus*.

107. *Cryptorrhynchus undulatus*, sp. n. (Tab. XXXIV. fig. 2, ♀.)

Oblong-ovate, shining, piceous or nigro-piceous, the antennæ and the tips of the tarsi ferruginous; somewhat thickly clothed with moderately coarse fulvous scales, with a few blackish scales intermixed, the elytra with a spot or short oblique streak on the fifth interstice below the base, and a narrow, angulate, postmedian fascia (of variable extent, and sometimes reduced to a few spots), white; the upper surface also set with coarse, suberect, blackish and pallid setæ, which are clustered into two small fascicles at the apex of the prothorax, as well as along the two slightly swollen portions of the third elytral interstice; the vestiture of the under surface sparse and pale, the legs setulose and more or less distinctly annulate. Head rugulosely punctate, the eyes rather prominent and well-separated; rostrum stout, curved, very little longer than the prothorax, slightly widened at the base, rugosely punctate and carinate, becoming smoother towards the tip, that of the ♀ finely punctate beyond the middle, the antennæ inserted at about midway, joint 2 of the funiculus a little shorter than 1, the club ovate, annulate, acuminate. Prothorax much broader than long, abruptly narrowed and constricted in front; densely punctate. Scutellum small. Elytra much wider than the prothorax, subparallel at the base, broadly produced at the apex; punctate-striate, the interstices punctulate and sometimes granulate, more or less convex, 3 slightly swollen below the base and towards the middle, and 5 also swollen below the base, the dorsal striæ thus appearing sinuate. Beneath coarsely punctate. Mesosternum much raised, very prominent, arcuate-emarginate. Femora feebly unidentate.

Length $3\frac{2}{3}$ – $4\frac{1}{2}$, breadth $1\frac{7}{8}$ – $2\frac{1}{3}$ millim. (♂ ♀.)

Hab. GUATEMALA, Cubilguitz, Senahu, and Panima in Vera Paz (*Champion*); PANAMA, Bugaba, Volcan de Chiriqui (*Champion*).

Numerous examples, both from Vera Paz and Chiriqui, varying in the extent of the angulate white fascia of the elytra, and in the number and length of the setæ. The peculiar white markings of the elytra are characteristic.

108. *Cryptorrhynchus decorus*, sp. n. (Tab. XXXIV. figg. 3, 3 a, ♀.)

Oblong-ovate, shining, nigro-piceous, the antennæ and the tips of the tarsi ferruginous; thickly clothed with rather coarse intermixed fulvous, whitish, and black scales—the black scales condensed into two short curved vittæ on the disc of the prothorax and various spots and streaks on the elytra, and the whitish scales into a narrow oblique humeral fascia; the upper surface also set with coarse, suberect, scattered setæ, which are clustered into two small fascicles at the apex of the prothorax and one on each of the swollen portions of the third elytral interstice, the legs setulose and subannulate. Head rugulosely punctate, the eyes rather narrowly separated; rostrum curved, not much longer than the prothorax, widened and rugulosely punctate at the base, and somewhat closely punctate thence to the apex, smoother along the middle, the antennæ inserted at about midway, joint 3–7 of the funiculus short,