

punctate-striate, the interstices punctulate and feebly convex. Beneath densely, finely punctate. Mesosternum raised, prominent, arcuate-emarginate. Legs rather slender, the tarsi long; femora each with a fine, sharp tooth.

Length $4-4\frac{1}{2}$, breadth $1\frac{3}{4}-2$ millim. (σ φ .)

Hab. MEXICO, Cerro de Plumas (*Höge*); GUATEMALA, Sinanja in Vera Paz (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).

A single example from each locality, agreeing very well *inter se*. This insect resembles *C. sedulus*, but it has a relatively narrower prothorax and more slender legs, and the vestiture is not so coarse. This last-mentioned character separates *C. erraticus* from *Graphonotus cinctipennis* and others. The strongly transverse, bimaculate prothorax, the much shorter elytral setæ, and the simple hind tibiae of the male distinguish it from *C. uncipes*.

111. *Cryptorrhynchus anthonomoides*, sp. n. (Tab. XXXIV. fig. 6, φ .)

Oblong-ovate, shining, piceous, the antennæ and the tips of the tarsi ferruginous; variegated above with a dense clothing of small brown, black, and whitish scales, the black scales on the prothorax condensed into two short curved vittæ on the disc, and those on the elytra into an irregular, interrupted, oblique fascia before the middle, various scattered spots between this and the base, and a few other spots towards the apex, the whitish scales clustered on the prothorax into a faint median vitta (or a spot in front of the scutellum), an oblique streak on the disc anteriorly, and a common post-median fascia; the upper surface also with a few, very short, scattered, decumbent, setiform scales; the vestiture of the legs and under surface pale brown, the legs with whitish scales intermixed. Head rugulose punctate, the eyes rather prominent, distant; rostrum curved, about as long as the prothorax, thickly punctate, rugose and slightly widened at the base, a little longer and smoother in the φ , the antennæ inserted behind the middle, the funiculus rather stout, joint 2 shorter than 1, the club ovate. Prothorax transverse, rounded at the sides, narrowed and feebly constricted in front, the ocular lobes not prominent; densely, finely punctate. Scutellum rounded. Elytra wider than the prothorax, subparallel in their basal half; seriate-punctate, the interstices closely punctulate, becoming slightly convex towards the sides, the third a little swollen below the base. Beneath closely, finely punctate. Mesosternum prominent, arcuate-emarginate. Legs rather short, similar in the two sexes; femora feebly clavate, each with a short tooth.

Length $4\frac{1}{5}-5$, breadth $1\frac{9}{10}-2\frac{1}{10}$ millim. (σ φ .)

Hab. GUATEMALA, Dueñas (*Champion*).

Seven specimens, varying in the colour of the vestiture, the whitish markings not always distinct. Very like *Graphonotus cinctipennis* and *G. lituratus*, but with a much shorter rostrum and a prominent arcuate-emarginate mesosternum, the prothorax nigro-bivittate, the elytra more mottled with black. It is a little more elongate than *C. erraticus*; the upper surface is variegated with whitish or cinereous, and the prothoracic vittæ are longer and narrower.

112. *Cryptorrhynchus fulvofasciatus*, sp. n.

Oblong-ovate, shining, piceous, the legs and antennæ rufo-piceous; the head, the prothorax with the sides, flanks, and a narrow space down the middle, the scutellum, and the elytra with a rather broad, irregular, subapical fascia, a few spots across the middle of the disc, and some others at the apex, clothed with fulvous scales, the rest of the squamosity of the prothorax and elytra sparse and blackish, that of the legs and under surface fulvous, fine, and scattered; the upper surface also set with short, semierect setæ. Head densely punctate, the eyes large and rather narrowly separated; rostrum short, stout, curved,