

upper surface also thickly set with long, erect setæ, the legs setose. Head densely punctate, the eyes moderately large, distant; rostrum short, stout, feebly curved, widened at the base, thickly punctate, the apical half smoother in the ♀, the antennæ inserted at or a little behind the middle, the club ovate. Prothorax transverse, rounded at the sides, feebly constricted and much narrowed in front; densely punctate. Scutellum small. Elytra slightly wider than the prothorax, parallel at the base; punctate-striate, the interstices feebly convex. Beneath densely punctate. Mesosternum very prominent, raised, arcuate-emarginate. Legs short; femora sharply unidentate.

Length $3-3\frac{1}{8}$, breadth $1\frac{1}{2}$ millim. (♂ ♀.)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One pair. Near *C. sedulus*, but much smaller and with long setæ, the elytra with an interrupted black and white sutural stripe and scattered black and white spots, the prothorax with a white median vitta.

116. *Cryptorrhynchus unipunctatus*, sp. n. (Tab. XXXIV. fig. 9, ♀.)

Oblong-ovate, shining, nigro-piceous, the antennæ, the tip of the rostrum, and the tarsi ferruginous; thickly clothed with blackish-brown scales, the prothorax and elytra mottled with black, the scutellum and a transverse spot on the suture below the base white; the upper surface also set with numerous short, coarse, blackish, erect setæ, which are clustered into two small fascicles at the apex of the prothorax; the vestiture of the under surface sparse, cinereous, the legs setose, cinereo-annulate. Head rugulosely punctate, the eyes distant; rostrum feebly curved, about as long as the prothorax, rugosely punctate and a little widened at the base, and finely punctate thence to the tip, the antennæ inserted slightly behind the middle, the club ovate. Prothorax transverse, rounded at the sides, narrowed and constricted in front; densely punctate. Scutellum small. Elytra wider than the prothorax, subparallel at the base; seriate-punctate, the interstices punctulate and almost flat. Beneath closely punctate. Mesosternum very prominent, raised, v-shaped in front. Legs short; femora acutely unidentate.

Length $3\frac{1}{2}$, breadth $1\frac{2}{3}$ millim. (♀.)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Two specimens. In this insect the transverse white spot on the suture is very conspicuous, owing to the unusually dark vestiture of the rest of the elytral surface. The black markings are asymmetric.

117. *Cryptorrhynchus abjectus*, sp. n. (Tab. XXXIV. fig. 10, ♂.)

Subovate, shining, piceous, the elytra sometimes mottled with rufo-piceous, the antennæ and the tips of the tarsi ferruginous; thickly clothed with intermixed pale brown, white, and blackish scales, the white scales clustered into various small scattered spots on the disc of the elytra (and in one specimen into two faint curved lines on each side of the prothorax) and the dark ones into two small patches at the base of the prothorax, a streak on the third elytral interstice at the base, and some irregular markings beyond this; the upper surface also set with short, erect, blackish and pallid setæ, the blackish setæ becoming more crowded at the base of the third elytral interstice, the legs setulose. Head densely punctate, the eyes distant; rostrum stout, curved, not longer than the prothorax, slightly widened at the base, coarsely, closely punctate in the ♂, much smoother in the ♀, the antennæ inserted at the middle, joint 2 of the funiculus half the length of 1, the club ovate. Prothorax transverse, rounded at the sides, narrowed and feebly constricted in front, slightly sinuate at the base; closely punctate. Scutellum small, rounded. Elytra a little wider than the prothorax, parallel in their basal third, the humeri obtuse; punctate-striate, the interstices feebly convex, 3 slightly swollen at the base. Beneath densely punctate. Mesosternum arcuate-emarginate. Legs short; femora feebly unidentate.

Length $3\frac{1}{2}$, breadth $1\frac{2}{3}$ millim. (♂ ♀.)