

the base subtruncate; with rows of rather coarse, rounded punctures, the interstices densely punctulate and more or less convex. Beneath dull, closely, finely punctate. Mesosternum raised and prominent, v-shaped in front. Femora sharply unidentate. Tarsi rather long and slender.

Length $3-3\frac{3}{4}$, breadth $1\frac{2}{5}-1\frac{2}{3}$ millim. (σ φ .)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Twelve specimens, varying in the colour of the scales, but with the general markings constant. The double ring-like mark on the disc of the prothorax (suggestive of a pair of spectacles) is characteristic. The metathoracic episterna are broad.

120. *Cryptorrhynchus murinus*, sp. n. (Tab. XXXIV. figg. 13, 13 a.)

Oblong-ovate, narrow, piceous, the antennæ, the tip of the rostrum, and the apices of the tarsi ferruginous; variegated above with a dense clothing of white, brownish-cinereous, and blackish scales, with short, decumbent, coarser, setiform scales intermixed, the white scales (when present) clustered into an interrupted median line and several spots on the prothorax, and an oblique fascia and various small scattered spots on the elytra, the blackish scales on the latter usually condensed into an elongate streak on the suture at the base, a shorter streak below it, and various irregular spots; the vestiture of the legs and under surface brownish-cinereous or whitish, the legs fusco-annulate. Head densely punctate, the eyes very large and rather narrowly separated; rostrum curved, a little longer than the prothorax, rugulose punctate, carinate, and slightly widened at the base, for the rest sparsely, very finely punctate, the antennæ inserted near the middle in the σ , and at about one-third from the base in the φ , joints 1 and 2 of the funiculus subequal in length, 3-7 short, the club oblong-ovate. Prothorax transverse, narrowed and feebly constricted in front, bisinuate at the base, the ocular lobes not very prominent; densely, finely punctate. Scutellum small. Elytra a little wider than the prothorax, subparallel in their basal half, the humeri rather prominent; seriate-punctate, the interstices convex, rugulose. Beneath dull, densely punctate, the metasternum and first ventral segment broadly depressed in the middle in the σ . Mesosternum raised, prominent, v-shaped in front. Legs rather short; femora unidentate; tarsi narrow.

Length $2\frac{1}{2}-3\frac{1}{2}$, breadth $1-1\frac{1}{2}$ millim. (σ φ .)

Hab. PANAMA, Bugaba, Volcan de Chiriqui (*Champion*).

Numerous examples, four only of which are distinctly marked with white, the others (mostly more or less discoloured) having the vestiture brownish with darker markings.

121. *Cryptorrhynchus propinquus*, sp. n. (Tab. XXXIV. fig. 14, φ .)

Larger and a little more elongate than *C. murinus*; the vestiture reddish-brown, with a few white scales intermixed, the black scales very similarly distributed, the mesothoracic side-pieces usually with an ochreous patch; the eyes not quite so large, less approximate; the ocular lobes of the prothorax a little more prominent.

Length $3\frac{2}{5}-4\frac{4}{5}$, breadth $1\frac{1}{2}-2$ millim. (σ φ .)

Hab. GUATEMALA, Cerro Zunil, Teleman (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).

Six specimens, four of which are from Cerro Zunil, differing too much from the long series of the variable *C. murinus* to be included under that species.