

large, rather coarsely faceted, and narrowly separated; rostrum curved, a little longer than the prothorax, finely punctate, the basal half rugulose and feebly carinate, smoother in the ♀, the antennæ inserted at (♀) or slightly beyond (♂) the middle, joints 2-7 of the funiculus short, the club ovate. Prothorax transverse, rounded at the sides anteriorly, much narrowed in front; densely, finely punctate and obsoletely carinate. Scutellum rounded. Elytra rather short, considerably wider than the prothorax, narrowing from near the base, the humeri obtuse; finely punctate-striate, the interstices punctulate, 3, 5, 7, and 9 more or less raised, 3 slightly swollen before and beyond the middle, 9 subcostate. Beneath densely punctate. Mesosternum U-shaped. Femora feebly unidentate. Tibiæ almost straight, subangulate at the base externally.

Length  $3\frac{1}{5}$ – $3\frac{3}{5}$ , breadth  $1\frac{1}{2}$ – $1\frac{7}{8}$  millim. (♂ ♀.)

*Hab.* GUATEMALA, Chacoj in Vera Paz, El Tumbador, El Reposo (*Champion*).

Four specimens. A small, ovate form, with close blackish vestiture, a white scutellum, and two clusters of coarse, erect, velvety-black scales on each elytron near the suture, the posterior two sometimes united into a common, curved fascia. *C. quadrisignatus* might perhaps be included amongst the smaller forms here placed in Section B of *Eubulus*.

Species small, ovate, robust, thickly squamose, with the rostrum short and curved, the eyes very large, finely faceted, and narrowly separated, the antennal club oblong-ovate, the prothorax bisinuate at the base, with prominent ocular lobes, the scutellum small, the elytra oval, with concentric markings, sinuate at the sides beneath, the mesosternum prominent, feebly, subangularly emarginate, the metathoracic episterna moderately broad, the ventral segment 2 longer than 3, the legs short and stout, the femora unidentate. (No. 124.)

#### 124. *Cryptorrhynchus concentricus*, sp. n. (Tab. XXXIV. fig. 17.)

Subovate, robust, black, the antennæ ferruginous; densely clothed with fulvous scales, the prothorax with coarser scales, which become sparser and darker on the disc and paler or whitish across the anterior constriction; the elytra with a streak along the suture behind the scutellum, a common annulus at the base, an interrupted curved line beyond it, and a faint transverse fascia near the apex, black, the annulus incompletely bordered within and without by a more or less distinct whitish line, the apex and a space in front of the dark line sometimes slightly suffused with whitish; the upper surface also set with very short, decumbent, setiform scales; the vestiture of the under surface close and pale, that of the legs fulvous, the femora cinereous at the base. Head rugulosely punctate, the eyes very large, finely faceted, rather narrowly separated; rostrum short, curved, rugosely punctate, obsoletely carinate at the base, the apical half more sparsely punctate in the ♀, the antennæ inserted at the middle, joint 2 of the funiculus a little shorter than 1, 3-7 transverse, the club oblong-ovate. Prothorax transverse, abruptly narrowed and strongly constricted in front; densely punctate. Scutellum small, rounded. Elytra considerably wider than the prothorax, narrowing from a little below the base, the humeri rounded; seriate-punctate, the interstices densely punctulate, feebly convex. Beneath opaque, densely punctate. Mesosternum raised, very prominent, v-shaped. Legs short and stout; femora sharply unidentate.

Length  $4\frac{1}{2}$ –5, breadth  $2\frac{1}{3}$ – $2\frac{1}{2}$  millim. (♂ ♀.)

*Hab.* PANAMA, Bugaba (*Champion*).

Three specimens. The peculiar elytral markings readily distinguish *C. concentricus*. It is rather like some of the species here placed under *Tyrannion*, but has very large, finely-faceted eyes.