

Species moderately large, subovate, densely squamose and setulose, with the eyes prominent, angular above, the rostrum very stout, short, and arcuate, the antennal club oblong-ovate, the prothorax deeply bisinuate at the base and fasciculate on the disc, the elytra subtriangular, with the alternate interstices sharply costate, the mesosternum arcuate-emarginate, the ventral segments 2-4 equal in length, the femora unidentate. (No. 125.)

125. *Cryptorrhynchus discedens*, sp. n. (Tab. XXXIV. figg. 18, 18 *a*, ♂.)

Subovate, black, the antennæ ferruginous; thickly clothed with rather coarse fulvous or rufo-fulvous scales, the upper surface slightly mottled with ochreous or whitish (the elytra in one specimen with intermixed blackish scales), and also set with short, scattered, semierect, setiform scales, which are clustered into six fascicles on the prothorax, the legs variegated with lighter and darker scales. Head rugosely punctate, irregularly, transversely carinate or tuberculate in front, the eyes rather prominent, well-separated, angular above; rostrum very stout, short, curved, coarsely, rugosely punctate, swollen at the tip in the ♂, the apical half smoother in the ♀, the antennæ inserted beyond the middle, joints 3-7 of the funiculus moniliform, 2 shorter than 1, the club stout, oblong-ovate, compact. Prothorax transverse, abruptly narrowed and constricted in front; densely punctate, with four setigerous prominences placed in a transverse series across the middle and two at the apex. Scutellum small, smooth. Elytra somewhat triangular, much wider than the prothorax, the humeri swollen and rather prominent; seriate-punctate, the interstices 3, 5, 7, and 9 sharply costate. Beneath coarsely, closely punctate; mesosternum prominent, arcuate-emarginate. Legs rather short; femora unidentate.

Length 5-6½, breadth 2½-3⅔ millim. (♂ ♀.)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Two specimens, the female larger and more robust than the male, with the vestiture coarser and more variegated, and the femoral teeth larger and blunter. This species has somewhat the facies of *Oxypterus asper* (Boh.). The very short rostrum, which is swollen at the tip in the male, separates it from *Eubulus*.

Species small, subovate, thickly squamose and setulose, with the rostrum curved, widened towards the base, the antennal club oblong-ovate or ovate, the eyes very large, finely faceted, and narrowly separated, the prothorax bisinuate at the base, the elytra subtriangular, with the alternate interstices not or moderately raised, the mesosternum very prominent, v-shaped in front, the metathoracic episterna broad, the ventral segment 2 longer than 3, the femora clavate and sharply unidentate, the anterior pair stouter than the others. (Nos. 126-130.)

126. *Cryptorrhynchus scutulatus*, sp. n. (Tab. XXXIV. figg. 19, 19 *a*, ♂.)

Subovate, nigro-piceous, the antennæ ferruginous; variegated with a dense clothing of intermixed fulvous and cinereous scales, the prothorax and elytra here and there mottled or interruptedly reticulate with black, the elytra with a dense fulvous spot on each shoulder in front, the legs fusco-annulate; the upper surface also sparsely set with short, erect, setiform scales, which are clustered into four small fascicles on the prothorax, two on the disc and two at the apex; the vestiture of the under surface sparse, cinereous. Head densely punctate, transversely flattened above the very large, narrowly separated eyes; rostrum strongly curved, a little longer than the prothorax, widened, rugulose, and squamose at the base, and for the rest closely, finely punctate, except along the smooth median line, the antennæ inserted slightly behind the middle, joints 1 and 2 of the funiculus subequal in length, the club oblong-ovate. Prothorax transverse, somewhat rounded at the sides, narrowed and feebly constricted in front; densely punctate, feebly binodose on the disc and at the apex. Elytra wider than the prothorax, rather short, narrowed from a little below the base; seriate-punctate, the alternate interstices a little raised. Beneath