

dull, finely punctate. Mesosternum raised, very prominent, v-shaped in front. Femora more or less clavate, acutely unidentate. Posterior tibiae slightly hollowed at the apex within.
Length $3-3\frac{1}{2}$, breadth $1\frac{1}{2}-1\frac{3}{4}$ millim. (σ .)

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Four specimens, probably all males. This insect has the general facies of *Graphonotus variisquamis*, but it is much smaller, the black markings of the upper surface are reduced to a few scattered spots or short streaks, the prothorax is feebly binodose on the disc and at the apex, and the prominent mesosternum is v-shaped in front. The eyes are more approximate than in *C. propinquus*, the humeri more prominent, &c.

127. Cryptorrhynchus pictipes, sp. n. (Tab. XXXIV. figg. 20, 20 a, ♀.)

Oblong-ovate, nigro-piceous, the antennae ferruginous; thickly clothed with fulvous scales, with a few cinereous scales intermixed, the prothorax spotted and the elytra interruptedly reticulate with black, the latter also sparsely set with small suberect scales; the legs nigro-annulate. Head densely punctate, the eyes very large and subcontiguous; rostrum rather slender, strongly curved, a little longer than the prothorax, widened, rugulose, and subcarinate at the base, for the rest shining and almost smooth, the antennae inserted at about the basal third, the club oblong-ovate, about as long as the preceding six joints united, joints 1 and 2 of the funiculus subequal in length. Prothorax broader than long, rounded at the sides anteriorly, narrowed and constricted in front, deeply bisinuate at the base; densely punctate, smoother at the apex. Elytra a little wider than the prothorax, gradually narrowing from about the basal fourth; seriate-punctate, the interstices (1 and 2 excepted) convex, 3, 5, and 7 more raised than the others. Beneath closely punctate. Mesosternum prominent, raised, v-shaped in front. Femora sharply unidentate, the anterior pair strongly, the others more feebly, clavate.

Length 4, breadth $1\frac{4}{5}$ millim. (♀.)

Hab. PANAMA, Bugaba (*Champion*).

One specimen. More elongate than *C. scutulatus*, the prothorax less transverse, and without fascicles of erect setae on the disc (the latter, however, may be abraded), the anterior femora strongly clavate.

128. Cryptorrhynchus decoratus, sp. n.

Oblong-ovate, shining, rufo-piceous, variegated with black, the antennae ferruginous; thickly clothed with intermixed fulvous, black, and whitish scales, the whitish scales mostly confined to the flanks of the prothorax, a line down the middle of its disc, and a small scutellar patch, the black scales condensed into a short streak on the disc of the elytra at the base and various irregular markings below this; the upper surface also set with short, semierect, setiform scales, which are clustered into four small fascicles on the prothorax—two on the disc and two at the apex; the legs subannulate. Head densely punctate, the eyes oval and well-separated; rostrum curved, slightly longer than the prothorax, widened, rugulose, and subcarinate at the base, and for the rest finely punctate, the antennae inserted at the middle, joint 2 of the funiculus shorter than 1, the club oblong-ovate. Prothorax a little broader than long, rounded at the sides anteriorly, narrowed and feebly constricted in front; densely punctate, feebly binodose on the disc and at the apex. Elytra considerably wider than the prothorax, parallel at the base, the humeri rather prominent; seriate-punctate, the interstices feebly convex. Beneath finely punctate. Mesosternum prominent, raised, v-shaped in front. Legs rather long; femora acutely unidentate, the anterior pair rather strongly, and the others more feebly, clavate, the posterior pair reaching the apex of the elytra; posterior tibiae straight on their inner edge; tarsi slender.

Length $3\frac{1}{3}$, breadth $1\frac{1}{2}$ millim. (σ ?)

Hab. PANAMA, Bugaba (*Champion*).