

One specimen. Narrower than *C. scutulatus*, the eyes not so large, the prothorax with a whitish median line, which is continued along each side of the scutellum, the black markings on the elytra mostly included within a large, common, subtriangular space on the basal half of the disc, the alternate interstices not raised, the legs longer and more slender.

129. **Cryptorrhynchus erubescens**, sp. n. (Tab. XXXIV. figg. 21, 21 a.)

Subovate, nigro-piceous, the rostrum and the legs in part rufo-piceous, the antennæ ferruginous; variegated with a dense clothing of fulvous and rosy-red scales, the prothorax and elytra narrowly, irregularly reticulate with black, the upper surface also somewhat thickly set with erect setiform scales, which are clustered into four fascicles on the prothorax (two on the disc and two at the apex), the legs subannulate. Head densely punctate and shallowly canaliculate, with a large, rounded, flattened space above, which is limited by an oblique carina on each side near the very large, rather narrowly separated, finely faceted eyes; rostrum strongly curved, a little longer than the prothorax, widened, rugulose, and subcarinate at the base, and for the rest shining and minutely punctate, the antennæ inserted at about the basal third, the club stout and elongate, nearly as long as the funiculus. Prothorax transverse, subquadrate, compressed at the sides anteriorly, abruptly narrowed and feebly constricted at the apex, the apex emarginate in the middle and somewhat produced, the base deeply bisinuate; densely punctate, binodose on the disc and in front. Scutellum small. Elytra much wider than the prothorax, transversely convex, rounded-triangular, abruptly declivous at the sides; seriate-punctate, the interstices convex, 3, 5, and 8 considerably raised, 3 swollen at the middle. Beneath densely punctate. Mesosternum very prominent, raised, v-shaped in front. Legs short and stout; femora clavate, sharply unidentate.

Length $4\frac{1}{4}$, breadth $2\frac{1}{5}$ millim. (♀?)

Hab. NICARAGUA, Chontales (*Belt*).

One specimen. This interesting little insect has much the facies of a *Zygops*. The circularly flattened front, the large eyes, the subquadrate, laterally compressed, quadrifasciculate prothorax, the subtriangular elytra, with raised interstices, &c., give it a peculiar facies. The vestiture of the upper surface and legs is in part rosy-red.

130. **Cryptorrhynchus honestus**, sp. n. (Tab. XXXIV. figg. 22, 22 a, ♂.)

Subovate, rather narrow, dull, rufo-piceous, variegated with black, the antennæ, and the legs in great part, ferruginous; mottled above with a rather sparse clothing of small ochreous, white, and blackish scales, the dark portions of the surface (including a transverse patch on the head and a transverse angulate line on the middle of the disc of the prothorax) almost bare, the elytra with a white spot on the suture before the middle; the upper surface also set with short, semierect setæ, the prothorax with two fascicles of longer, blackish, erect setæ at the apex; the legs subannulate with small ochreous scales, the vestiture of the under surface sparse and pale. Head rugulosely punctate, the eyes large and narrowly separated; rostrum curved, not longer than the prothorax, closely, finely punctate, rugulose and a little widened at the base, the antennæ inserted at the middle, joint 2 of the funiculus shorter than 1, 3-7 short, the club ovate. Prothorax rather broader than long, subconical, narrowing from near the base, the latter deeply bisinuate; closely, finely punctate, binodose at the apex. Scutellum small. Elytra much wider than the prothorax, subtriangular; seriate-punctate, the interstices opaque, 3 and 5 slightly raised anteriorly, the others somewhat convex. Beneath finely punctate, opaque. Mesosternum very prominent, and much raised, v-shaped in front. Legs short, the anterior pair longer; femora feebly clavate, sharply unidentate.

Length 3, breadth $1\frac{2}{5}$ millim. (♂.)

Hab. PANAMA, Bugaba (*Champion*).