

rather deeply punctate-striate, the anterior femora obsoletely, and the others sharply, dentate. The elytra are a little shorter than in *C. echinatus*.

142. **Cryptorrhynchus pallescens**, sp. n.

Ovate, narrow, shining, rufo-piceous; clothed with pallid scales, which on the prothorax become a little more condensed along the middle and at the sides, the upper surface and legs also somewhat thickly set with long, erect, pallid setæ. Head densely punctate, the eyes distant; rostrum short, curved, rather sparsely punctate, rugulose at the base, the antennæ inserted at the middle, the club ovate. Prothorax transverse, feebly constricted and much narrowed in front, bisinuate at the base; densely punctate. Scutellum very small. Elytra a little wider than the prothorax, oval; rather coarsely punctate-striate, the interstices feebly convex. Beneath closely punctate. Mesosternum prominent, feebly emarginate. Legs short; femora each with a small tooth.

Length $2\frac{2}{3}$, breadth $1\frac{1}{5}$ millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One specimen. In this insect the vestiture is entirely pale. The prothorax appears to be faintly bivittate on the disc, but this is due to the scales being more crowded on the other portions of its surface. Near *C. pallidisetis*, but longer and larger, the setæ coarser, the elytra more coarsely punctate-striate. Smaller and more ovate than *C. tectus*, the setæ much longer, the prothorax more sinuate at the base and without black vittæ on the disc.

Species minute, oblong-ovate, densely squamose, and with a long, erect, abundant, pallid setosity, the rostrum short, straight, and flattened, the eyes rather small, the antennal club ovate, the prothorax bisinuate at the base, the scutellum minute, the elytra a little wider than the prothorax, the mesosternum arcuate-emarginate, the metathoracic episterna moderately broad, the ventral segment 2 slightly longer than 3, the legs short, the femora clavate, unidentate. (No. 143.)

143. **Cryptorrhynchus pallidisetis**, sp. n. (Tab. XXXV. fig. 3.)

Oblong-ovate, piceous, the rostrum, antennæ, and legs more or less ferruginous; densely clothed with brownish-white scales, the elytra mottled with pale brown, the upper surface also thickly set with very long, erect, fine, pallid setæ, the legs setose. Head densely, very finely punctate, the eyes rather small, distant, exposed; rostrum straight, flattened, about as long as the prothorax, very finely punctate, squamose at the base, the antennæ inserted behind the middle, the club ovate. Prothorax broader than long, narrowing from near the base, the base itself feebly sinuate; densely punctate. Scutellum small. Elytra very little wider than the prothorax, subparallel at the base; finely punctate-striate, the interstices almost flat. Beneath densely punctate. Mesosternum arcuate-emarginate. Legs very short; femora stout, sharply unidentate.

Length 2, breadth 1 millim.

Hab. MEXICO, Atoyac in Vera Cruz (*H. H. Smith*).

One specimen, somewhat immature, but with the vestiture in perfect condition.