

- b*¹. Posterior knees with the outer lobe not longer than the inner; prothorax strongly bisinuate at the base.
- a*². Prothorax with a large red patch on each side in front; posterior femora not annulate.
- a*³. Posterior femora bidentate; eyes well-separated; prothorax nigro-bivittate and nigro-fasciate; pygidium black . . . *rufomaculata*, sp. n.
- b*³. Posterior femora unidentate; eyes subcontiguous; prothorax with a dark patch on the disc; pygidium red . . . *erythropyga*, sp. n.
- b*². Prothorax without red markings; posterior femora sharply nigro-annulate.
- c*³. Posterior femora bidentate; posterior tibiae feebly mucronate at the inner apical angle.
- a*⁴. Prothorax with an anchor-shaped blackish patch on the disc, the sides albo-maculate posteriorly; elytra variegate . . . *mexicana*, Boh.
- b*⁴. Prothorax with a rounded or subquadrate dark patch on the disc, the sides obliquely streaked with white posteriorly; elytra with a black patch, preceded and followed by a white spot . . . *histrion*, Boh.
- d*³. Posterior femora unidentate; posterior tibiae flavo-annulate, sharply mucronate at the inner apical angle; prothorax with a subtriangular dark patch on the disc, nigro-maculate on the flanks; elytra variegate . . . *maculiventris*, sp. n.
- c*. Posterior femora shorter, not or about reaching the apex of the abdomen.
- c*¹. Posterior knees with the outer lobe broadly produced; posterior femora 4- or 5-dentate; body subelliptic: length over 9 mm. . . *quadridentata*, sp. n.
- d*¹. Posterior knees with the outer lobe not longer than the inner; body oblong or oblong-ovate: length $3\frac{1}{2}$ – $7\frac{1}{5}$ mm.
- c*³. Posterior femora 3- or 4-, the other femora 2-, dentate.
- e*³. Elytra with large fulvous, white-edged spots, the prothorax with a transverse, arcuate, black fascia; the vestiture of the under surface almost wholly white . . . *amænula*, sp. n.
- f*³. Elytra variegate, the prothorax not transversely fasciate; the vestiture of the under surface more or less mottled at the sides.
- c*⁴. Prothorax and elytra blackish, sprinkled with white . . . *disjecta*, sp. n.
- d*⁴. Prothorax and elytra variegate; posterior femora fuscous at the middle.
- a*⁵. Prothorax interruptedly trivittate; joint 3 of the funiculus as long as 4–7 united . . . *sellata*, sp. n.
- b*⁵. Prothorax spotted; joint 3 of the funiculus much shorter than 4–7 united . . . *tridentata*, Gyll.