

- a.* Elytra separately produced at the apex, the apices truncate or dentiform and often raised.
- a*¹. Species larger, with moderately long legs. [PARATIMORUS, Heller.]
- a*². Elytral vestiture coarse and overlapping, the apices not produced into a sharp spine.
- a*³. Elytra with scattered clusters of raised scales, forming tuberculiform prominences, spotted; prothorax fulvous at the sides . . . *ganglbaueri*, Heller.
- b*³. Elytra without tuberculiform prominences, arcuato-fasciate, the interstices here and there slightly swollen and setose; prothorax not fulvous at the sides . . . *triangulifer*, sp. n.
- c*³. Elytra flattened, without raised scales or setæ, the suture and two oblique radiating lines on each side of it fulvous or white; prothorax in great part fulvous . . . *biradiatus*, sp. n.
- b*². Elytral vestiture sparse and fine, the apices produced into a sharp point . . . *bicaudatus*, sp. n.
- b*¹. Species smaller, with shorter legs; elytral vestiture coarse and overlapping. [CYLINDROCOPTURUS, Heller.]
- c*². Second ventral segment with two conspicuous conical prominences*.
- d*³. Prothorax with a cluster of erect fulvous scales in front, the sides broadly white; elytra each with a triangular black patch on the disc, the sides angularly dilated before the apex . . . *tetralobus*, sp. n.
- e*³. Prothorax without erect scales in front, the sides with a broad white patch behind; elytra with a common \wedge -shaped pallid line below the base, the sides not dilated near the apex . . . *armatus*, sp. n.
- d*². Second ventral segment with the prominences indistinct or wanting.
- f*³. Prothorax narrower than the elytra.
- a*⁴. Prothorax with a large white patch at the sides behind; elytra with a common \wedge -shaped pallid line below the base, the apices dentiform . . . *imbricatus*, sp. n.
- b*⁴. Prothorax and elytra spotted with white, the elytra each with a transverse fulvous patch below the base, the apices obliquely truncate . . . *albonotatus*, sp. n.
- c*⁴. Prothorax interruptedly albo-trivittate, the elytra with two curved series of white spots, the apices truncate . . . *adpersus*, Lec.
- g*³. Prothorax and elytra subequal in width; prothorax albo-vittate on each side, the elytra bifasciate . . . *bifasciatus*, sp. n.
- b.* Elytra conjointly rounded at the apex, the interstices not wider than the striæ.
- c*¹. Upper surface with large scales; elytra each with a transverse or oblique black patch: larger, body elongate-elliptic . . . *elongatus*, sp. n.
- d*¹. Upper surface with smaller scales; prothorax with a large pallid patch at each angle; elytra mottled: smaller, body oblong-elliptic . . . *scaphiformis*, sp. n.

* Possibly a sexual character.