

Meso- and metasternum flattened and unimpressed between the widely-separated intermediate coxæ. Legs short and rather slender; femora unidentate, the posterior pair obsoletely carinate externally, the area on the anterior pair rather large, concave, and finely strigose.

Length 3, breadth  $1\frac{1}{2}$  millim.

*Hab.* GUATEMALA, Cerro Zunil (*Champion*).

One specimen. Smaller and less robust than *C. stridulans*, the eyes almost contiguous, the rostrum finely punctate to the tip, the legs more slender. The cinereous scales on the elytra appear to be a little more crowded at the middle of the first and second interstices.

### 8. *Copturomimus brevicollis*, sp. n.

Subovate, nigro-piceous, the elytra mottled with rufo-piceous, their apical margin, the rostrum, antennæ, and legs ferruginous; the vestiture fine and rather sparse: the head between the eyes and the prothorax ochreous, the latter with two bare spaces on the disc; the elytra mottled with ochreous and darker scales, the ochreous scales condensed into a small transverse patch on the suture beyond the middle and a narrow streak behind the scutellum; the under surface and legs sparsely clothed with narrow yellowish scales. Eyes very large, narrowly separated above. Rostrum moderately stout, not reaching beyond the middle coxæ, rugulose and carinate at the base. Prothorax nearly twice as broad as long, rounded at the sides, narrowed and constricted in front; densely, rugulosely punctate, and finely carinate. Elytra deeply punctate-striate, the interstices flat and rugulose. Meso- and metasternum flattened between the middle coxæ. Legs short and rather slender; femora unidentate, the posterior pair carinate externally, the area on the anterior pair rather large and finely strigose.

Length  $2\frac{3}{5}$ , breadth  $1\frac{1}{2}$  millim.

*Hab.* PANAMA, Volcan de Chiriqui (*Champion*).

One specimen. Very near the Guatemalan *C. sparsus*, but with the pallid scales finer, ochreous in colour, and clustered into definite markings, the rostrum not so stout, the prothorax shorter and more rounded at the sides, the eyes less approximate above.

### 9. *Copturomimus cinereus*. (Tab. IV. figg. 24; 24 a.)

*Copturomimus cinereus*, Heller, Abhandl. Mus. Dresd. no. 11, p. 64<sup>1</sup>.

*Hab.* MEXICO, Orizaba<sup>1</sup>, Toxpam (*Sallé*), Jalapa, Cerro de Plumas (*Höge*); PANAMA, Volcan de Chiriqui (*Champion*).

Six specimens of this species are before me, including one of the types. *C. cinereus* agrees with the Colombian *C. ochrofasciatus*, Heller (a co-type of which I have seen), in having the obliquely strigose area on the anterior femora small and partly hidden by the vestiture. The metasternal depression is rather deep.

### 10. *Copturomimus confluens*, sp. n. (Tab. IV. figg. 25, 25 a, b.)

Subovate, black, the prothorax and elytra sometimes piceous, the antennæ, rostrum, and tarsi more or less ferruginous; the vestiture rather fine above, coarse beneath: the prothorax with the sides broadly, and a space along the middle, ochreous; the elytra with a rather broad basal, median, and apical fascia (extending outwards to the sixth stria), connected along the suture, and a transverse patch behind the