

1. *Copturomorpha interrupta*, sp. n. (Tab. IV. fig. 26.)

Subovate, piceous, the rostrum and antennæ ferruginous; the prothorax sparsely clothed with rather coarse, narrow, ochreous scales, which are somewhat condensed on the middle of the disc, and with some coarser whitish scales at each hind angle; the elytra mottled with narrow ochreous and fuscous scales, and with a whitish sutural stripe, which is dilated below the base into a cruciform patch and interrupted at the middle by a common V-shaped fuscous mark, the seventh interstice with an ochreous spot at one-third from the apex and the fourth with a similar spot at the base; the vestiture of the under surface coarse, close, and whitish, the ventral segments 3-5 with a space down the middle, and 1 at the sides, almost bare, the legs with narrow whitish scales. Eyes very large, subcontiguous, the intra-ocular space gradually widening above and below. Rostrum rugulose and carinate at the extreme base. Joint 2 of the funiculus slightly longer than 1. Joint 1 of the antennal club longer than the others united. Prothorax transverse, narrowing from near the base, feebly constricted in front; finely, rugulose punctate, and feebly carinate. Elytra somewhat rounded at the sides, blunt at the tip, depressed along the suture; deeply striate, the striæ finely punctate, the interstices flat and rugulose. Mesosternum broadly depressed, the depression limited on each side by an oblique ridge. Prosternum with a conical prominence behind each coxa, and the coxa itself with a similar prominence. Legs short; femora unarmed beneath and without external carina, the area on the anterior pair very finely strigose, the intermediate and posterior knees with an acute tooth.

Length $4\frac{3}{4}$, breadth $2\frac{1}{2}$ millim.

Hab. MEXICO, Jalapa (*Smith, ex H. F. Wickham*).

One specimen, presented by Mr. Wickham. This insect has the general facies of *Copturomimus stridulans*; but the femora are unarmed beneath, the posterior knees are spinose, and the mesosternum is shaped as in the genus *Pæcilogaster*.

2. *Copturomorpha 11-notata*, sp. n. (Tab. IV. figg. 27, 27 a, b.)

Subovate, piceous; somewhat thickly clothed with piliform ochreous scales, the prothorax with a patch at each hind angle, a short streak on the basal lobe, and the flanks, the elytra with a small transverse patch on the suture at about one-third from the base, a short streak on the suture at one-third from the tip, a minute spot at the base of the fourth interstice, another at the base of the seventh, and yet another on the seventh at one-third from the apex, and the under surface, clothed with rather coarse white scales. Eyes very large, subcontiguous, separated above by a narrow lanciform space. Rostrum rugulose and carinate at the base. Funiculus with joints 1 and 2 rather elongate, subequal in length, 3 and 4 much shorter, 5-7 very short or transverse. Prothorax transverse, somewhat rounded at the sides, narrowed and constricted in front; densely, finely punctate, and feebly carinate. Elytra punctate-striate, the interstices flat and rugulose. Meso- and metasternum broadly depressed between the middle coxæ, the mesosternal depression limited on each side by a fine oblique ridge. Prosternum with a dentiform prominence, similar to that on each anterior coxa. Legs short; femora unarmed beneath, the finely strigose area on the anterior pair just traceable, the posterior knees with an acute tooth.

Length $3-3\frac{1}{2}$, breadth $1\frac{1}{2}-2$ millim.

Hab. GUATEMALA, San Gerónimo (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).

Two specimens, the one from Guatemala immature and partly abraded, the other in perfect condition. Smaller than *C. interrupta*, the two basal joints of the funiculus subequal in length, the markings different.