

3. *Copturomorpha rugipennis*, sp. n.

Subovate, nigro-piceous, the antennæ, the tip of the rostrum, and the tarsi more or less ferruginous; sparsely clothed with narrow pale ochreous scales, with darker scales intermixed: the ochreous scales on the prothorax mostly clustered along the sides, at the middle of the base, and in the depressed space in front, and those on the elytra into a transverse basal fascia, a streak on the suture behind the scutellum and another at the middle, and a spot at the sides; the under surface with coarser, oval, pallid scales, the vestiture of the legs similarly-coloured, the posterior femora with an oblique dark ring. Eyes subcontiguous. Rostrum rugulose and finely carinate at the base. Joints 1-2 of the funiculus equal in length, 3 a little shorter, 4-7 short or transverse. Prothorax short, rather convex, constricted and narrowed in front; densely, rugosely punctate, and feebly carinate. Elytra much wider than the prothorax; deeply punctate-striate, the interstices narrow, convex, rugose. Mesosternum deeply excavate between the coxæ. Legs short; femora unarmed beneath, the finely strigose area on the anterior pair inconspicuous and placed near the base.

Length $3\frac{1}{3}$, breadth $1\frac{3}{4}$ millim.

Hab. MEXICO, Toxpam in Vera Cruz (*Sallé*).

One specimen. In this species the scattered vestiture is in great part ochreous above, and the incomplete sutural stripe is abruptly interrupted before the middle by a transverse dark patch. Near *C. 11-notata*, but with the elytral sculpture coarser, and the transverse white patch on the suture replaced by a darker one.

4. *Copturomorpha albomaculata*, sp. n. (Tab. IV. fig. 28.)

Subovate, black, the antennæ, tip of the rostrum, and tarsi ferruginous; variegated above with oval white and small narrow black scales: the white scales on the prothorax clustered into three large spots at the base, another in the centre in front, and a minute spot at each anterior angle, and those on the elytra into a narrow sutural stripe, which is broadly interrupted beyond the middle, a short streak at the base of each dorsal interstice, another near the middle of the second, a spot on the seventh at about one-third from the apex, and sometimes another on the fourth in a line with it, the rest of the elytral surface with black scales; the under surface and legs clothed with white scales. Head bare above; eyes subcontiguous; rostrum carinate and rugulose at the base; joints 1 and 2 of the funiculus equal in length. Prothorax short, constricted and narrowed in front; densely, rugulosely punctate, and feebly carinate. Elytra rounded at the sides, blunt at the apex; deeply punctate-striate, the interstices about as wide as the striæ, flattened. Mesosternum deeply excavate between the coxæ, the metasternum abruptly declivous in front. Legs short; femora sulcate and unarmed beneath, the strigose area on the anterior pair inconspicuous and placed near the base.

Length $2\frac{2}{5}$ - $2\frac{3}{5}$, breadth $1\frac{1}{4}$ - $1\frac{1}{2}$ millim.

Hab. GUATEMALA, Dueñas (*Champion*).

Three specimens. Narrower than *C. albosignata* and *C. leucosticta*, the mesosternum deeply excavate, the prothorax with a minute white spot at each anterior angle, the elytra with the white markings less extended, &c. The white scales are coarse, above and beneath.

5. *Copturomorpha albosignata*, sp. n. (Tab. IV. figg. 29, 29 a.)

Subovate, piceous or black, the antennæ, the tips of the tarsi, and the apical margin of the elytra ferruginous; the vestiture close, coarse, long, and piliform: the prothorax with three large patches at the base (the outer one extending forward along the flanks beneath) and a broad transverse band in front, and the elytra with a common cruciform patch at the base, a transverse band below the shoulder,