

and a common, broad, undulate, subapical fascia, white, the rest of the elytral vestiture black or blackish, that of the under surface and legs wholly white. Head bare above, closely punctate, without smooth space on the vertex, foveate between the eyes, which are moderately large and narrowly separated; rostrum sparsely punctate, rugulose and carinate at the base, smooth along the middle; joints 1 and 2 of the funiculus equal in length, 3 and 4 much shorter, also equal in length. Prothorax short, rather convex, constricted and narrowed in front; densely, finely punctate, and also carinate. Elytra rather short and broad, depressed along the suture, bluntly and separately rounded at the apex; shining, deeply punctate-striate, the interstices convex, rugulosely punctate. Meso- and metasternum slightly depressed between the middle coxæ. Legs short; femora unarmed beneath and at the apex, and without external carina, the strigose area on the anterior pair oblong.

Var. The elytra with the common subapical fascia reduced to three spots, the transverse subhumeral mark wanting.

Length $2\frac{3}{4}$ – $3\frac{1}{2}$, breadth $1\frac{1}{2}$ –2 millim.

Hab. GUATEMALA, Cerro Zunil (*Champion*); PANAMA, Volcan de Chiriqui (*Champion*).

Three specimens of each form, one only from Guatemala. In this species the markings on the prothorax and elytra are clothed with long, coarse, narrow, hair-like scales, a character separating *C. albosignata* from its nearest allies.

6. *Copturomorpha leucosticta*, sp. n. (Tab. IV. fig. 30.)

Subovate, piceous, the antennæ and tarsi, and the apical margin of the elytra, ferruginous; clothed with oval white scales, the upper surface with intermixed small, brown or fulvous, hair-like scales: the white scales on the prothorax condensed into a median vitta, a transverse patch on each side at the apex, and a large patch at each hind angle (the lateral markings connected externally and almost enclosing subtriangular bare spot), and those on the elytra into a broad sutural stripe (which is interrupted before the middle), a spot on the shoulder, another at the side towards the apex, and a complete or much interrupted line on each dorsal interstice; the legs with narrow white scales. Head bare above, closely punctate; eyes large, subcontiguous in front; rostrum rugulose and subcarinate at the base; joints 1 and 2 of the funiculus equal in length. Prothorax short, densely, rugosely punctate. Elytra shining, deeply punctate-striate, the interstices not wider than the striæ, rugulose and subgranulate. Mesosternum slightly depressed between the coxæ. Legs short; femora unarmed, the strigose area on the anterior pair small.

Length 3 – $3\frac{1}{3}$, breadth $1\frac{1}{2}$ – $1\frac{2}{3}$ millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

Two specimens, one with the dorsal interstices of the elytra albolineate to near the apex, the other with the white lines reduced to a few scattered streaks. Near *C. albosignata*, but with the white scales on the upper and under surfaces oval in shape, the head not foveate between the eyes, the elytra more rugose.

7. *Copturomorpha funerea*, sp. n.

Short ovate, black, the antennæ, the apical margin of the elytra, the apices of the tibiæ, and the tarsi, ferruginous; the prothorax with a few narrow white scales at the hind angles and along the sides; the elytra clothed with narrow black and white scales, the latter condensed into some small spots at the base, a spot at the middle of the suture, another at its apex, and another on the eighth interstice; the vestiture of the legs and under surface white, the posterior femora with an oblique fuscous band beyond the middle. Eyes almost contiguous in front, narrowly separated above. Rostrum thickly punctate to the tip, rugose at the base. Joint 2 of the funiculus a little longer than 1. Prothorax very short, rounded at the sides, constricted and narrowed in front; densely, rugosely punctate, and obsoletely carinate. Elytra rather