

29. *Baris punctirostris*, sp. n.

Oblong-ovate, very finely alutaceous, moderately shining, æneous, the tibiae sometimes reddish; the punctures at the sides of the prothorax, along the elytral interstices, and on the under surface and legs each bearing a very small, setiform scale. Head closely, minutely punctate, transversely grooved between the eyes; rostrum very short, stout, arcuate, closely punctate to the tip in both sexes. Prothorax transverse, gradually narrowing from the base, constricted in front; densely punctate, except along the smooth median line. Scutellum small, rounded, concave. Elytra sharply, narrowly striate, the striae feebly punctate, the first deeply impressed from a little below the base, the interstices broad, flat, closely uniseriate-punctate, the third, and sometimes the second also, confusedly punctured, the ninth raised towards the apex. Beneath closely punctate; metasternum and first ventral segment hollowed in the middle in the ♂. Tibiae almost straight.

Length $3-3\frac{1}{2}$, breadth $1\frac{1}{3}-1\frac{1}{2}$ millim. (♂ ♀.)

Hab. GUATEMALA, Quezaltenango 7800 feet (*Champion*).

Three specimens. An oblong, æneous form very near *B. ærea*, with a closely punctured rostrum, a densely punctured prothorax (except along the sharply-defined median line), and closely seriate-punctate, finely setulose elytra. More elongate than *B. setosella*, the elytral interstices less densely and not so coarsely punctate. More closely punctate above (the rostrum included) than *B. corrusca*, the elytral vestiture conspicuous.

30. *Baris seriatosetosa*. (Tab. XXII. figg. 23, 23 a, ♂.)

Baris seriatosetosa, Solari, Ann. Mus. Genova, xlii. p. 419 (1906) ¹.

Hab. MEXICO, "Sierra de Durango" ¹.

A small, oblong-ovate, somewhat depressed form, æneo-piceous in colour, with an extremely short rostrum, a densely punctured prothorax (without trace of smooth median line), a minute, subquadrate, smooth scutellum, and closely, finely uniseriate-punctate elytral interstices, the punctures on the elytra each bearing a small, conspicuous, setiform scale. *B. seriatosetosa* comes very near some of the varieties of *B. ærea*.

31. *Baris macraspis*, sp. n. (Tab. XXII. figg. 24, 24 a, ♀.)

Subovate, shining, piceous, with a faint æneous tinge; glabrous above, the punctures on the under surface and legs each bearing a minute scale. Head very sparsely, minutely punctate, transversely grooved between the eyes; rostrum stout, strongly arcuate, shorter than the prothorax, flattened at the tip, sparsely, minutely punctate. Prothorax transverse, compressed at the sides in front, sparsely, somewhat finely punctate. Scutellum narrowed posteriorly, oblong, small, flattened. Elytra a little wider than the prothorax, rounded-subtriangular, depressed along the suture, narrowly striate, the interstices very broad, flat, each with a row of extremely minute scattered punctures, the ninth raised towards the apex. Beneath closely, the abdomen sparsely and finely, punctate. Prosternum slightly depressed along the middle to near the base. Tibiae straight.

Length $3\frac{1}{2}$, breadth $1\frac{1}{5}$ millim. (♀.)

Hab. GUATEMALA, Dueñas (*Champion*).

One specimen. A subovate form, with a sparsely punctured prothorax, very broad,