

♂ ♀. Similar to the following genus (*Piezocorynus*), but the legs rough with a long pubescence, first tarsal segment shorter, second broader.

- a.* Pronotal carina angulate in middle, the angle pointing forwards . . . . . Species No. 2.  
     Pronotal carina not angulate in middle, or the angle pointing backwards . . . . . *b.*  
*b.* Suture of elytra dotted with black . . . . . Species No. 1.  
     Suture of elytra with one large spot before middle . . . . . Species No. 3.

### 1. *Lagopezus tenuicornis*.

*Anthribus tenuicornis*, Fabr. Syst. Eleuth. ii. p. 407 (1801)<sup>1</sup>; Schoenh. Gen. Spec. Curc. i. p. 182. n. 7 (1833)<sup>2</sup>.

*Lagopezus tenuicornis*, Fähr. in Schoenh. Gen. Spec. Curc. v. p. 190. n. 1 (1839)<sup>3</sup>; Labr. & Imh. Gen. Curc. i. no. 51 (1842)<sup>4</sup>; Lacord. Gen. Col. vii. p. 541 (1866)<sup>5</sup>; Gemm. & Har. Cat. Col. ix. p. 2736 (1872)<sup>6</sup>.

*Hab.* NICARAGUA, Chontales (*Belt, Janson*); PANAMA, Bugaba, Volcan de Chiriqui (*Champion*). — SOUTH AMERICA<sup>1</sup>, Colombia<sup>5</sup>, Cayenne<sup>4 5 6</sup>, Peru<sup>5</sup>, Upper Amazons, Pernambuco, Pará, Espiritu Santo (*Mus. Tring*).

Found in some numbers in Nicaragua and Panama.

### 2. *Lagopezus inversus*, sp. n. (Tab. X. fig. 23.)

♂. In colour similar to *L. tenuicornis*, Fabr. (1801), but the suture of elytra not tessellated. Proportionally shorter; upperside more irregularly grooved; frons broader; tenth antennal segment more than twice as broad as long; dorsal carina triflexuose, forming in middle an obtuse but very distinct angle which points forward; basal callosity of elytron higher and more abrupt.

Length 6 millim.

*Hab.* GUATEMALA, Cubilguitz in Vera Paz (*Champion*).

One male.

### 3. *Lagopezus morio*, sp. n. (Tab. X. fig. 24.)

In general appearance not unlike *Piezocorynus tristis*, Jekel (1855).

♂ ♀. Black, covered with a greyish-black pubescence which is somewhat paler than in *L. tenuicornis*, with the following velvety-black markings: on pronotum two oblique apical lines converging anteriorly, a basal patch on each side more or less produced beyond carina; on elytra a sutural streak from near base to middle, the suture dotted with black from this streak to apex; antenna rufescent, last segment luteous. Rostrum shorter than in *L. tenuicornis*; frons wider; pronotum shorter, with the basal carina rather farther from the base in middle, the mesial vestigial callosity standing close to the carina, the side less incurved before basal angle, the lateral carina being much less flexed; elytra shorter, interspaces more even, with traces of tubercles, which are more or less tipped with black, basal callosity higher than in *L. tenuicornis*, more abrupt at sutural side.

Length 5 millim.

*Hab.* PANAMA, Volcan de Chiriqui, Bugaba, Tolé (*Champion*).

Two males, one female.