

8. *Piezocorynus simplex*, sp. n. (Tab. XI. fig. 3.)

♂. Similar to the preceding. Rostrum shorter. Antennæ much thinner, segment 8 broader, 9 a little longer than 8, 10 hardly twice as broad as long. Pronotum with apical mesial streak as in the preceding species, disc visibly depressed on each side of the slightly raised mesial line, also more depressed before carina than in *P. strigifer*; carina very feebly concave in middle, lateral angle less rounded off than in *P. strigifer*. Alternate interspaces of elytron subcostate, first three interspaces nearly as in *P. strigifer*, fifth also dotted with luteous. First three abdominal segments flattened.

Length 3 millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

One male.

9. *Piezocorynus plagifer*. (Tab. X. figg. 25, 25 a.)

Piezocorynus dispar, Lec. (nec Gyllenhal, 1833), Proc. Amer. Philos. Soc. xv. p. 402. n. 1 (1876)¹.

Piezocorynus plagifer, Jordan, Nov. Zool. xi. p. 277. n. 71 (1904)².

Hab. NORTH AMERICA, Missouri and Texas¹.—GUATEMALA, Cerro Zunil (*Champion*), Coban in Vera Paz (*Conradt*); NICARAGUA, Chontales (*Belt*); COSTA RICA²; PANAMA, Bugaba, Volcan de Chiriqui (*Champion*).—BRAZIL².

Now I have seen more specimens of *P. plagifer* I come to the conclusion that it may possibly intergrade with *P. dispar*, Gyll., being perhaps only a colour-variety of that insect.

Found in some numbers in Central America.

XI. ALLANDRUS.

Allandrus, Leconte, Proc. Amer. Philos. Soc. xv. p. 396 (1876) (type: *bifasciatus*); Jord. Nov. Zool. xi. p. 242 (1904).

Very close to *Enedreytes*, Schoenh. (1839).

Range: America.

1. *Allandrus angulatus*, sp. n. (Tab. XI. fig. 4.)

♂ ♀. Black, variegated with grey; antenna (except club), tibiae, and tarsi more or less rufous. Rostrum flat, not carinate, densely rugate-punctate, about half as long again as apically broad. Eye elliptical. Prothorax rugate-punctate, with a luteous patch before scutellum; carina convex, angulate in middle, the point of the angle directed forwards. Scutellum luteous. Elytra broader than in *A. bifasciatus*, Lec., from North America, the grey pubescence a little denser before and behind, forming indistinct and irregular transverse bands.

Length 2½ millim.

Hab. PANAMA, Volcan de Chiriqui 2500–4000 feet (*Champion*).

A small series.