

genus. *Eugonus* can be divided into two groups, the species of the typical group having the sterna smooth at the sides, the second group having them punctured laterally. In Central America only two species, belonging to the second group, have so far been found.

Suture and lateral edge of elytra dotted Species No. 1.
Suture and lateral edge of elytra not dotted Species No. 2.

1. *Eugonus subcylindricus*.

Eugonus subcylindricus, Fåhræus, in Schoenh. Gen. Spec. Curc. v. p. 172. n. 2 (1839)¹; Gemm. & Har. Cat. Col. ix. p. 2743 (1872)².

Schimotocheilus subcylindricus, Fåhræus (Chevrolat, in litt.), l. c. (sub syn.)³.

Schimotocheilus bostrichoides, Fåhræus (ex Dejean, Cat.), l. c. (sub syn.)⁴.

Eugonus fallax, Gemm. & Har. (ex Dejean), l. c. (sub syn.)⁵.

(?) *Phænisor albofasciatus*, Motsch. Bull. Soc. Imp. Nat. Moscou, xlviii. 2, p. 241 (1874)⁶.

♂ ♀. A rather common species of wide distribution and great individual variability. The pubescence of the elytra is generally buffish-ochraceous, but sometimes whitish-grey.

Hab. MEXICO^{1 2 3}, Toxpam, Playa Vicente (*Sallé*); GUATEMALA, Yzabal (*Sallé*), San Juan, Chacoj, and Panzos in Vera Paz (*Champion*); NICARAGUA, Chontales (*Belt, Janson*).—BRAZIL^{4 5}.

2. *Eugonus decorus*, sp. n. (Tab. XIII. fig. 10.)

♂. Close to *E. ornatus*, Jord. (1904), from Brazil, the ninth antennal segment longer, the pattern different, and the legs not rufous except the knee-joints and the claws. Pronotum with a large but not sharply marked patch on each side extending forward beyond middle, touching the second of two small black dorso-lateral spots standing between apical margin and middle. Elytra greyish-white, ornamented with the following black markings: a round spot on shoulder; a cruciform sutural patch at base; a transverse submedian band consisting of an elliptical sutural patch and a larger lateral patch which is sinuate at the lateral margin and behind and is connected with the basal patch behind the shoulder; a large round subapical patch which reaches neither suture nor outer margin.

Length 5 millim.

Hab. NICARAGUA, Chontales (*Janson*).

One male. Perhaps only a geographical race of *E. ornatus*, of which we have a series.

XXIII. EUGONODES.

Eugonodes, Jordan, Nov. Zool. xi. p. 302 (1904) (type: *marmoreus*).

Like *Eugonus*, but with the apical sinus of rostrum vestigial or absent.

1. *Eugonodes brevirostris*.

♀. *Eugonodes brevirostris*, Jordan, l. c. n. 130 (1904)¹.

A single ♂, which has small crests of hairs on the abdomen.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).—BRAZIL, Bahia¹ (*Mus. Tring*).