

One male. More shining than its allies, the prothorax with a broader smooth space on the middle of the disc, the ventral excavation ( $\sigma$ ) deep and with a transverse prominence in the centre behind, thus appearing arcuate in shape.

## 2. *Tytthomimus rufotestaceus*, sp. n.

Oblong-ovate, narrow, moderately shining, faintly alutaceous, ferruginous or rufo-testaceous. Head almost smooth, transversely depressed and sometimes foveate between the eyes, the latter moderately large and separated by less than their own width; rostrum short, stout, feebly curved, parallel-sided in its outer half, sparsely, very finely punctate, the antennæ inserted at the middle. Prothorax not quite so long as broad, suboval, the subapical constriction indistinct, obliterated above; closely, finely punctate, with an abbreviated smooth median line at the base. Scutellum almost smooth, conspicuous. Elytra moderately long, a little wider than the prothorax, subparallel in their basal half, broadly produced at the apex, the apices somewhat explanate and bluntly, separately rounded; narrowly punctate-striate, the interstices broader than the striæ, convex at the apex, finely uniseriate-punctate. Beneath sparsely, finely punctate; first ventral segment broadly depressed down the middle anteriorly in the  $\sigma$ . Tibiæ feebly mucronate at the inner apical angle, the anterior pair sinuate within.

Length  $2-2\frac{1}{10}$ , breadth  $\frac{2}{3}-\frac{3}{4}$  millim. ( $\sigma$  ♀.)

*Hab.* GUATEMALA, Trece Aguas in Alta Vera Paz (*Barber & Schwarz, in U.S. Nat. Mus.*:  $\sigma$ ), Las Mercedes, Pantaleon (*Champion*: ♀).

One specimen from each locality. Smaller and duller than *T. rubicundus*; the impunctate space on the disc of the prothorax reduced to a narrow line at the base, the subapical constriction almost wanting; the ventral depression of the male shallow and not extending to the apex. The relatively longer elytra, the smaller prothorax, the longer and more slender legs, &c., separate *T. rufotestaceus* from *Caulophilus sericatus*.

## 3. *Tytthomimus lineatocollis*, sp. n.

Moderately elongate, narrow, somewhat depressed, obscure ferruginous, finely alutaceous, feebly shining. Head closely, finely punctate on the vertex, foveate and slightly depressed between the eyes, the latter large and separated by a little more than half the width of the rostrum; rostrum cylindrical, rather slender, about as long as the posterior portion of the prothorax, sparsely, finely punctate, the antennæ inserted at the middle. Prothorax nearly as long as broad, somewhat oval, abruptly constricted in front; closely, finely punctate, except along the smooth median line, which extends from the base to the subapical constriction. Scutellum small. Elytra moderately long, slightly wider than the prothorax, subparallel to beyond the middle, obtusely produced and subexplanate at the apex; narrowly punctate-striate, the interstices much broader than the striæ, flat, transversely rugulose, and closely, conspicuously uniseriate-punctate. Beneath closely, finely punctate; first ventral segment narrowly depressed down the middle. Tibiæ each with a minute tooth at the inner apical angle, the anterior pair angularly dilated at the middle within.

Length  $2\frac{1}{4}$ , breadth  $\frac{2}{3}$  millim. ( $\sigma$ .)

*Hab.* PANAMA, Bugaba (*Champion*).

One specimen. More elongate and more depressed than the two other species referred to this genus, the head punctate on the vertex, the rostrum longer and not so stout, the prothorax with an almost complete impunctate median line, the elytral interstices transversely rugulose and conspicuously uniseriate-punctate.