

- g*<sup>2</sup>. Prothorax finely, subuniformly punctate, except along a narrow smooth space down the middle, even; elytra sharply punctate-striate; anterior tibiae more or less produced at the outer apical angle: upper surface almost bare.
- e*<sup>3</sup>. Punctuation of the prothorax sparser; elytral interstices very finely punctate . . . . . *memnonius*, Gyll.
- f*<sup>3</sup>. Punctuation of the prothorax a little coarser and closer; elytral interstices coarsely uniseriate-punctate . . . . . *vomerinus*, Lec.
- h*<sup>1</sup>. Rostrum feebly compressed or subcylindrical, more slender, not or scarcely widened towards the apex beneath (except in *S. incurrens*).
- h*<sup>2</sup>. Prothorax with a V-shaped area at the base, an oblong subapical depression, and a similar depression near each hind angle, coarsely punctate, the rest of the disc to near the sides smooth or very finely punctured; elytra rather coarsely punctate-striate: surface in great part bare, the punctures sometimes incrustate.
- g*<sup>3</sup>. Upper surface shining, black . . . . . *incurrens*, Gyll.
- h*<sup>3</sup>. Upper surface opaque, varying in colour from black to ferruginous; elytra often with the striae more coarsely punctate . . . . . *incurrens*, v. *monilis*, [Gyll.]
- i*<sup>2</sup>. Prothorax with a V-shaped or oblong, coarsely punctured, flattened or depressed, area at the base, the rest of the surface more or less finely punctate, smooth along the median line; elytra crenato- or punctato-sulcate: surface bare and shining.
- i*<sup>3</sup>. Body ferruginous or rufo-castaneous above, black beneath; the prothorax with three sharply-defined vittae on the disc, the scutellum, and the elytra sometimes with the alternate interstices, black; elytra crenato-striate.
- a*<sup>4</sup>. Fifth ventral segment sharply hollowed down the middle posteriorly, and the anterior and intermediate femora ciliate, in ♂ . . . . . *quadrivittatus*, Gyll.
- b*<sup>4</sup>. Fifth ventral segment without definite groove, and the anterior and intermediate femora penicillate at the base, in ♂ . . . . . *lineatus*, sp. n.
- j*<sup>3</sup>. Body black (the prothorax and elytra sometimes rufo-lineate or castaneous); elytra punctato-sulcate, with flat, finely punctate interstices . . . . . *imus*, Gyll.
- j*<sup>2</sup>. Prothorax narrow, with a small space in front of the scutellum rather more coarsely punctured than the rest of the minutely punctate surface, smooth along the median line; elytra sharply punctate-striate, the interstices flat, minutely punctate; anterior tibiae produced at their outer apical angle: surface bare and shining . . . . . *angustus*, Boh.