

APPENDIX TO THE CURCULIONINÆ.

SINCE the conclusion of Vol. IV. pts. 4 and 5, and of the earlier portion of pt. 7 (pp. 1-78), of this series*, various additional species belonging to the Subfamily Curculioninæ have come to hand, including a representative of the group "Gymnetrina," from Northern Mexico, and several new *Hilipus*, *Anthonomus*, *Cholus*, *Erethistes*, *Cratosomus*, &c., from more southern localities, sufficient to make a short Appendix desirable. Corrections to the synonymy (some of which have already been mentioned in scattered footnotes, &c.), and additional localities for several species (a few of which had been described from single examples), are also noted.

Vol. IV. pt. 4.

HYLOBIINA.

HILIPUS (pp. 7, 715).

Hilipus cruciatus (p. 9). (Tab. IX. fig. 2, var.)

Amongst the specimens of this species in the British Museum there is one from "Mexico" with the white markings on the elytra reduced to three oblique fasciæ and a transverse spot below the shoulder.

Hilipus fenestratus (p. 10).

To the locality given, add:—MEXICO, Colonia (*Flohr, in coll. Fry*).

Hilipus limbatus (p. 11).

To the locality given, add:—MEXICO (*ex coll. Solari*).

We are indebted to Signor Solari for a Mexican representative of this species, described from a single example from Chiriqui. He also has *H. albopictus* from Mexico, which again was described from a single specimen.

* Part 6 is devoted to the Brentidæ, Scolytidæ, and Anthribidæ.