

## ISCHIOMASTUS.

*Ischiomastus*, Kirsch, Abhandl. Zool. Mus. Dresden, 1888-89, no. 4, p. 27; Faust, Stett. ent. Zeit. 1893, p. 6.

*Lixomorpha*, Champ. antea, iv. 4, p. 62 (1902).

This correction in the synonymy has already been briefly noted on p. 1 of the present volume.

*I. marginatus* was omitted from our enumeration of the Hylobiina.

1. *Ischiomastus albomarginatus*.

*Lixomorpha albomarginata*, Champ. antea, iv. 4, pp. 63, 719.

2. *Ischiomastus marginatus*.

*Ischiomastus marginatus*, Faust. Stett. ent. Zeit. 1893, p. 10<sup>1</sup>.

*Hab.* COSTA RICA [♂]<sup>1</sup>.—COLOMBIA, Cauca Valley [♀]<sup>1</sup>.

Dr. Heller has kindly communicated one of the types (the ♀) of this species for examination. It is allied to *I. albomarginatus*.

## ANCHONINA.

## DIOPTROPHORUS (p. 93).

Amongst some *Dioptraphori* recently sent us by Mr. Wickham, from Tres Marias (Morelos) \* and Salazar, Mexico, there is one specimen (from Tres Marias) perhaps referable to *D. ocularis*, the type of which cannot now be found in the "Sommer" collection. It is a little larger and broader than *D. simplex*, and has the prothorax more uneven, more widened forwards, and abruptly constricted anteriorly.

## CLEONINA.

## LIXUS (p. 101).

15 (A). *Lixus breviculus*, sp. n.

♂. Comparatively short, rather narrow, black, the antennæ (joints 1-3 of the club excepted) and tarsi obscure ferruginous; sparsely clothed with very fine cinereous pubescence. Head and rostrum densely, finely punctate, the latter curved, stout, and a little shorter than the prothorax, the eyes somewhat prominent, the head feebly foveate between them. Prothorax strongly transverse, gradually narrowing from the base; densely, minutely punctate, with numerous coarser punctures intermixed, the disc slightly depressed down the middle at the base and apex. Elytra very little wider than the prothorax, moderately long, rounded at the tip; seriate-punctate, the interstices flat, densely punctulate, the scutellar region transversely depressed, the disc also with a shallow rounded foveiform depression at about one-third from the apex. Beneath densely, finely punctate, with a few coarser punctures intermixed; first ventral segment shallowly sulcate down the middle. Legs comparatively slender, the anterior femora unarmed. Wings (apparently) wanting.

Length 8, breadth  $2\frac{7}{8}$  millim.

\* Mr. Wickham has also sent us an *Anchonus* very nearly related to *A. granulatus* from this locality. It may represent yet another species of the genus.