

1. **Tristachycera viridis.** (Tab. III. fig. 7.)

Tristachycera viridis, Bates, loc. cit. p. 171.

Hab. NICARAGUA, Chontales (*Belt*).

ACHRYSON.

Achryson, Serv. Ann. Soc. Ent. Fr. 1833, p. 572.

Exclusively American, extending a little beyond the tropics, both north and south. Eleven species have been described.

1. **Achryson surinamum.**

Cerambyx surinamus, Linn. Syst. Nat. i. p. 632 (1767).

Cerambyx surinamensis, Oliv. Ent. no. 67, p. 42, t. 13. f. 93 (1795).

Stenocorus circumflexus, Fab. Mant. Ins. i. p. 144 (1787).

Var. *chontalense*, Bates, Trans. Ent. Soc. 1872, p. 171.

Hab. NORTH AMERICA, Middle States.—MEXICO; GUATEMALA; NICARAGUA; COSTA RICA.—SOUTH AMERICA, generally distributed to Paraguay.

HAMMATICHERUS.

Hammaticherus, Serv. Ann. Soc. Ent. Fr. 1834, p. 15.

Exclusively American, but closely allied to *Plocæderus*, which is an African and Indian genus. Eight species of *Hammaticherus* are known.

1. **Hammaticherus castaneus.** (Tab III. fig. 2.)

Hammaticherus castaneus, Bates, Trans. Ent. Soc. 1870, p. 250.

Hab. NICARAGUA, Chontales (*Belt, Janson*).—SOUTH BRAZIL.

2. **Hammaticherus mexicanus.**

Hammaticherus mexicanus, Thoms. Class. Long. p. 196; Bates, Trans. Ent. Soc. 1872, p. 171.

Hab. MEXICO.

ATYLOSTAGMA.

Atylostagma, White, Cat. Long. Col. B. M. i. p. 97.

The following is the sole known species:—

1. **Atylostagma politum.**

Atylostagma politum, White, loc. cit. p. 97, t. 3. f. 5.

Hab. HONDURAS (*Dyson*).

SPHALLENUM.

Sphallenum, Bates, Trans. Ent. Soc. 1870, p. 253.

Six species, exclusively Tropical-American.