

1. **Odontocera monostigma.** (Tab. V. fig. 6.)*Agaone monostigma*, Bates, Trans. Ent. Soc. 1869, p. 384.*Odontocera monostigma*, Bates, Ann. & Mag. Nat. Hist. xi. 1873, p. 38.*Hab.* NICARAGUA, Chontales (*Belt*).2. **Odontocera clara.** (Tab. V. fig. 4.)*Odontocera clara*, Bates, Ann. & Mag. Nat. Hist. xi. 1873, p. 38.*Hab.* NICARAGUA, Chontales (*Bates*).3. **Odontocera aurocincta.***Odontocera aurocincta*, Bates, Ann. & Mag. Nat. Hist. xi. 1873, p. 42.*Hab.* MEXICO, Tehuantepec (*Sumichrast*); Plan del Rio (*Höge*).4. **Odontocera rugicollis.**

O. bisulcatae primo intuitu similis, at differt inter alia femoribus posticis subito clavatis. Nigra; thorace fere cylindrico, dorso sulcis duabus haud profundis longitudinalibus, omnino grosse medio transversim rugulose punctato, disco rufo-castaneo; antennis brevibus, robustis, nigris; elytris modice abbreviatis, a medio dehiscentibus, apicem versus extrorsum paullo curvatis apice ipso obtuse truncatis, marginibus utrinque nigris punctatis, vitta mediana alba a basi usque ad apicem vitrea; corpore subtus nigro, episternis rufo-castaneis; pedibus nigris, femoribus tibiisque posticis basi albis, tibiis apicem versus paullulum sensim dilatatis extus nigro-pilosis.

Long. $6\frac{1}{2}$ lin.*Hab.* GUATEMALA, Zapote (*Champion*).

Much less elongated than *O. sanguinolenta*, and similar in form and colours to *O. bisulcata*; but distinguished from both by the abruptly clavate hind femora, which are nearly of the same form (relatively shorter) as in *O. gracilis*. Each elytron forms, after the middle, a linear parallel-sided blade, slightly flexuous, the two blades being more distant from each other near the middle than at the apex.

ACYPHODERES.

Acyphoderes, Serville, Ann. Soc. Ent. Fr. 1833, p. 549.

About a dozen species of *Acyphoderes* are known, having a similar distribution to the *Odontocerae* and *Ommatae*.

1. **Acyphoderes acutipennis.***Acyphoderes acutipennis*, Thoms. Class. Long. p. 179.*Hab.* MEXICO.2. **Acyphoderes vespiventris.**

Angustus, gracilis, nigro-chalybeus, antennis tibiis et tarsis piceis, partibus oris fulvo-rufis, elytris chalybeo-violaceis politis; thorace antice dilatato, dorso convexo postice trinodoso, alveolato-punctato, argenteo-piloso; scutello argenteo; elytris abdomine paullo brevioribus, angustis, fere linearibus, apice breviter truncatis, humeris elevatis, basi et apice aspere punctatis; antennis dimidiam corporis longitudinem fere