

COREMIA*.

Coremia, Serville, Ann. Soc. Ent. Fr. 1834, p. 22.

Three species are recorded from Tropical South America, one of which is found to extend its range to Central America.

1. *Coremia hirtipes*.

Saperda hirtipes, Oliv. Ent. no. 68, p. 14, t. 1. f. 8¹.

Coremia hirtipes, Guér. Icon. Règne Anim. t. 44. f. 3 b²; Bates, Trans. Ent. Soc. 1872, p. 186³.

Hab. NICARAGUA, Chontales (*Belt*³).—SOUTH AMERICA, Cayenne¹, Amazons.

Nicaraguan examples agree perfectly with those from the Amazons, but they are a little more robust and of a darker general tint.

CYLLENE.

Cyllene, Newman, Entom. p. 7 (1840).

Clytus, Serville, Laporte & Gory, White.

This well-marked group of the cosmopolitan *Clytini* comprises upwards of twenty species, all American, ranging from Buenos Ayres to California and Canada, but most numerous in the tropical zone.

1. *Cyllene erythropus*.

Clytus erythropus, Chevr. Col. du Mexique, Cent. i. fasc. 4. no. 1 (1835); Laporte & Gory, Mon. Clyt. p. 4, t. 2. f. 1.

Clytus variegatus, Laporte & Gory, Mon. Clyt. t. 2. f. 4.

Cyllene erythropus, Chevr. Ann. Soc. Ent. Fr. 1860, p. 458.

Hab. MEXICO, Jalapa (*Höge*); GUATEMALA, Capetillo (*Champion*).

2. *Cyllene guttata*.

Cyllene guttatus, Chevr. Ann. Soc. Ent. Fr. 1860, p. 459¹; Bates, Trans. Ent. Soc. 1872, p. 186².

Hab. MEXICO, Cordova (*Sallé*)¹; NICARAGUA, Chontales (*Belt*², *Janson*); COSTA RICA, Irazu (*Rogers*).

3. *Cyllene crinicornis*.

Cyllene crinicornis, Chevr. Ann. Soc. Ent. Fr. 1860, p. 460¹.

Hab. MEXICO, Granada, Vera Cruz¹; HONDURAS, river Sarstoon (*Blancaneaux*).

4. *Cyllene mexicana*.

Clytus mexicanus, Laporte & Gory, Mon. Clyt. p. 3, t. 2. f. 6.

Hab. MEXICO.

* *Ethemon lepidum*, recorded in the Munich Catalogue as from Mexico, is a South-Brazilian insect.